



















104  
43  
71361-

2

THE

VOYAGE OF H.M.S. CHALLENGER.

ZOOLOGY—VOL. XXIV.

*PLATES.*

27





8 QL  
438  
G78  
1888  
INVZ

Great Britain: Challenger Office.

508.2  
C437  
Zoo. 1888

REPORT  
ON THE  
SCIENTIFIC RESULTS  
OF THE  
VOYAGE OF H.M.S. CHALLENGER  
DURING THE YEARS 1873-76

UNDER THE COMMAND OF  
CAPTAIN GEORGE S. NARES, R.N., F.R.S.  
AND THE LATE  
CAPTAIN FRANK TOURLE THOMSON, R.N.

PREPARED UNDER THE SUPERINTENDENCE OF  
THE LATE  
Sir C. WYVILLE THOMSON, Knt., F.R.S., &c.  
REGIUS PROFESSOR OF NATURAL HISTORY IN THE UNIVERSITY OF EDINBURGH  
DIRECTOR OF THE CIVILIAN SCIENTIFIC STAFF ON BOARD

AND NOW OF  
JOHN MURRAY  
ONE OF THE NATURALISTS OF THE EXPEDITION

ZOOLOGY—VOL. XXIV.  
PLATES

c. 194973

Published by Order of Her Majesty's Government



PRINTED FOR HER MAJESTY'S STATIONERY OFFICE  
AND SOLD BY  
LONDON :—EYRE & SPOTTISWOODE, EAST HARDING STREET, FETTER LANE  
EDINBURGH :—ADAM & CHARLES BLACK  
DUBLIN :—HODGES, FIGGIS, & CO.

1888

PRINTED BY NEILL AND COMPANY, EDINEURGH,  
FOR HER MAJESTY'S STATIONERY OFFICE.



## C O N T E N T S.

REPORT on the CRUSTACEA MACRURA dredged by H.M.S. CHALLENGER during the  
years 1873-1876.

By C. SPENCE BATE, F.R.S., &c.



## EDITORIAL NOTE.

THESE Plates illustrate the text of Mr. Spence Bate's Report on the CRUSTACEA MACRURA, and number in all 157—two of them illustrating Dr. Hock's Appendix to the Report on a parasitic CIRRIPED.

JOHN MURRAY.

CHALLENGER OFFICE, 32 QUEEN STREET,  
EDINBURGH, *8th May* 1888.





PLATE I.

## PLATE I.

---

The same letters represent homological parts on all the plates.

---

### *Cheiroplatea cenobita* (p. 12).

- Fig. 1♀. Lateral view; enlarged twice.  
.. 1*a*. Ophthalmopod or eye.  
.. 1*b*. First antenna.  
.. 1*c*. Second antenna.  
.. 1*d*. Siagon or mandible.  
.. 1*e*. First siagnopod or maxilla.  
.. 1*f*. Second siagnopod or maxilla.  
.. 1*g*. Third siagnopod or maxilla (injured ?).  
.. 1*h*. First gnathopod.  
.. 1*i*. Second gnathopod.  
.. 1*k*. First pair of pereiopoda, chelæ.  
.. 1*u*. Fourth pereiopod; terminal joints.  
.. 1*o*. Fifth pereiopod; terminal joints.  
.. 1*br*. Branchial plume.  
.. 1*r*. String of ova.  
.. 1*vzv*. Rhipidura; *z*, telson; *v,v*, posterior pair of pleopoda.

### *Cheramus orientalis* (p. 30).

- .. 2. Lateral view; enlarged four times.  
.. 2*c*. Cephalon and carapace (dorsal view).  
.. 2*a*. Ophthalmopod or eye.  
.. 2*b*. First antenna. Flagella.  
.. 2*i*. Second gnathopod.  
.. 2*o*. Fifth pereiopod; terminal joints.  
.. 2*q*. Second pleopod.  
.. 2*r*. Third pleopod.  
.. 2*s*. Fourth pleopod; *ms*, stylamblys, cincinnuli, and hair; magnified.  
.. 2*br*. Branchial plume.  
.. 2*br*. Section of branchia.  
.. 2*v*. Posterior pleopoda, margin; magnified.  
.. 2*z*. Telson.







PLATE II.

PLATE II.

*Cheramus occidentalis* (p. 32).

- Fig. 1. Lateral view; enlarged two and a half times.  
,, 1C. Cephalon, lateral view of carapace, with ophthalmopod, first antenna, and peduncle of second.  
.. 1Z. Telson; *v*, sixth pair of pleopoda.

*Callianassa occidentalis* (p. 29).

- ,, 2*k*. First pereopod; natural size.

*Scallasis amboinæ* (p. 34).

- ,, 3. Lateral view; enlarged four times.  
,, 3*a,a*. Ophthalmopoda.  
,, 3*b*. First antenna.  
,, 3*i*. Second pair of gnathopoda (attached).  
,, 3*o*. Fifth pereopod.  
,, 3*r*. Third pleopod.  
.. 3*r.ms*. Stylamblys; magnified.  
,, 3*br*. Branchial plume.  
.. 3*vzv*. Rhipidura; *z*, telson; *v,v*, posterior pair of pleopoda.  
.. 4. Ventral aspect; enlarged six times. Posterior somites of pleon and rhipidura reversed.







PLATE III.

PLATE III.

*Thalassina scorpionoides*, nat. size (p. 19).

- c.* Peduncle of first antenna; *c''*, coxal joint seen in section, showing the acoustic apparatus; *c'''*, arrangement of hairs within the chamber; *c''''*, one of the hairs magnified.
- d.* Siagon or mandible.
- e.* First siagnopod.
- f.* Second siagnopod, with one of the hairs of the mastigobranchial lash magnified in sections.



J.S. Van der Horst, J. Hawley

THALASSINA SCORPIONOIDES





PLATE IV.

*Thalassina scorpionoides*, branchiæ (p. 19).

Fig. 1. Pereion, showing the form and arrangement of the branchiæ; lateral wall of the carapace removed.

- „ *h*. First gnathopod with rudimentary mastigobranchia, with podobranchia and arthrobranchia attached; *h''*, arthrobranchial plume (detached); enlarged.
- „ *i*. Second gnathopod with rudimentary mastigobranchia, with podobranchia attached; *i''*, arthrobranchial plume detached, showing the under side; enlarged.
- „ *k*. Coxa of first pereiopod with rudimentary mastigobranchial lash, with the podobranchial plume attached; *k''*, anterior arthrobranchia detached, showing under side; *k'''*, posterior arthrobranchia, showing upper surface; *k'''*, under surface of the base of the same; enlarged.





1844

THALASSINA SCORPIONIDES, BRANCHIA



PLATE V.

PLATE V.

*Paraxius altus* (p. 37).

- Fig. 1. Lateral view; enlarged three times.  
,, 1. Rostrum, upper surface.  
.. 1*d*. Siagon or mandible.  
,, 1*g*. Third siagnopod.  
,, 1*h*. First gnathopod.  
,, 1*i*. Second gnathopod (with podobranchia and mastigobranchia detached)  
,, 1*k*. First pereopod.  
,, 1*l*. Second pereopod.  
,, 1*o*. Fifth pereopod (extremity).  
,, 1*q*. Second pleopod (part).  
,, 1*vzv*. Rhipidura; z, telson; *v,v*, pleopoda.  
.. 1*br*. Branchial plumes.

*Eiconaxius acutifrons* (p. 40).

- .. 2♀. Lateral view; enlarged three and a half times.  
.. 2♀. Upper surface of rostrum, female.  
.. 2♂. Rostrum, male.  
.. 2*d*. Siagon or mandible.  
.. 2*g*. Third siagnopod.  
.. 2*h*. First gnathopod.  
,, 2*i*. Second gnathopod.  
,, 2*kl*. First pereopod; from the left side.  
.. 2*q*. Third pleopod, with stylamblys and a cincinnulus detached and magnified.

*Eiconaxius kermadeci*, female (p. 43).

- .. 3♀. Lateral view of a female specimen; enlarged twice.  
,, 3*kl*. First pereopod; from the left side.  
,, 3*kr*. First pereopod; from the right side.  
,, 3*l*. Second pereopod.

*Eiconaxius parvus* (p. 44).

- ,, 4. Lateral view of a female specimen; enlarged four times.  
,, 5. Brephalos extracted from the ovum, seen from the ventral surface; magnified about thirty times.



1. PARAXIUS ALTUS  
2. EICONAXIUS ACUTIFRONS.  
3. EICONAXIUS KERMADECI  
4. EICONAXIUS PARVUS  
5. EICONAXIUS PARVUS BREPHALUS



PLATE VI.



PLATE VI.

*Thaumastocheles zaleuca*, nat. size (p. 47).

- c. Ventral aspect of the cephalon.
- c''. Frontal aspect of the same.
- b. First antenna.
- c. Second antenna.
- h. First gnathopod.
- i. Second gnathopod.
- k'. First pereopod, showing the propodos.
- k''. First pereopod, showing the articulation of the dactylos.
- l'''. First pereopod, showing the articulation of the ischium with the basis.
- v. Fourth pereopod; terminal joints.
- w. Fifth pereopod: terminal joints.
- p. First pleopod.
- q. Second pleopod.





PLATE VII.

PLATE VII.

*Thaumastocheles zaleuca* (p. 47).

- Fig. 1. Pereion ; lateral view, with part of carapace removed to show the arrangement of the branchiæ.
- „ 1*br*. Diagram of the vertical arrangement of a section of the branchial apparatus.
- „ 1*c*. First siagnopod.
- „ 1*f*. Second siagnopod.
- „ 1*g*. Third siagnopod.
- „ 1*h*. First gnathopod.

*Ibaccus verdi* (p. 58).

- „ 2. Dorsal aspect of the pereion with the carapace removed to show the branchiæ; left side with the branchiæ intact, right shown in section.
- „ 2*d*. Siagones or mandibles ; *ff*, shows the position of the extremities of the second pair of siagnopoda.
- „ 2*e*. First pair of siagnopoda.
- „ 2*f*. Second pair of siagnopoda.
- „ 2*g*. Third pair of siagnopoda.
- „ 2*k*. First pereiopod ; showing the coxa with mastigobranchia and podobranchia attached.
- „ *ple'*. Upper surface of the pyloric apparatus.
- „ *ple''*. Under surface of the pyloric apparatus.







PLATE VII.

PLATE VIII.

*Ibuccus cerdã*, nat. size (p. 58).

- $\text{♂}$ . Male, showing ventral aspect.
- $\text{♀}$ . Female, showing dorsal aspect.
- $a$ . Cephalon; dorsal aspect;  $b$ , first antennæ;  $c$ , second antennæ.
- $\sigma\text{5}$ . Fifth pereopod of the male.
- $\sigma\text{5}$ . Fifth pereopod of the female.
- $\mu\text{1}$ . First pleopod of the male.
- $\mu\text{2}$ . First pleopod of the female.
- $\eta\text{1}$ . Second pleopod of the male.
- $\eta\text{2}$ . Second pleopod of the female.





PLATE IV.

PLATE IX.

*Ibaccus brevipes*. (p. 62).

- Fig. 1.     Dorsal aspect of the animal, male ; nat. size.  
.. 1*q*.     Second somite of pleon.  
.. 1*q*.     Second pleopod.

*Ibaccus alticrenatus* (p. 63).

- .. 2♀.     Dorsal view of a female ; nat. size.

*Arctus sordidus* (p. 66).

- .. 3.     Dorsal view of a female ; nat. size.  
.. 3''.     One side of the ventral aspect of pereion ; enlarged.  
.. 3*q* ♂ . Second pleopod of the male.  
.. 3*q* ♀ . Second pleopod of the female.  
.. 3*r* ♂ . Third pleopod of the male.  
.. 3*r* ♀ . Third pleopod of the female.

*Arctus orientalis* (p. 68).

- .. 4.     Dorsal view of the female ; nat. size.  
.. 4''.     One side of the ventral aspect of pereion ; enlarged.  
.. 4*o* ♂ . Extremity of posterior pereopod of the male.  
.. 4*o* ♀ . Extremity of posterior pereopod of the female.








PLATE X.

PLATE X.

*Arctus tuberculatus* (p. 70).

- Fig. 1. Lateral view ; natural size.  
„ 2. Dorsal view ; enlarged three times.  
„ 1♂. Fifth pair of pereopoda of the male.  
„ 1♀. Fifth pair of pereopoda of the female.

*Arctus immaturus* (p. 71).

- „ 3. Dorsal aspect of a specimen ; enlarged four times.  
„ 3''. Ventral aspect of pereion.

*Arctus pygmæus* (p. 73).

- „ 4♀. Dorsal aspect of a female ; enlarged four and a half times.  
„ 4c. Second antenna.  
„ 4q. Third pleopod and stylamblys ; the latter detached and magnified.  
„ 4r. Brephalos ; extracted from the ovum.



1 ARCTUS TUBERCULATUS

2 " " ENLARGED

3 ARCTUS PYGMALUS

4 " PYGMALUS





PLATE Xa.

*Panulirus guttatus*, var., nat. size (p. 78).

c. Part of second antenna, showing the stridulating organ.







PLATE XL

PLATE XI.

*Palinostus lalandii* (p. 86).

(*Palinurus lalandii* on Plate.)

Fig 1. A young specimen ; twice the natural size.

„ 1*q*. Second pleopod.


*Panulirus angulatus* (p. 81).

„ 2. Ventral aspect of a young and imperfect specimen ; enlarged two and a half times.

„ 2*q*. Second pleopod.

„ 3. Dorsal aspect.

„ 4. Lateral aspect.



3. PANULIRUS ANGULATUS. DORSAL  
4. LATERAL



PLATE XI.

PLATE XI<sub>A</sub>.

*Palinostus lalandii*, nat. size (p. 86).







PLATE XII.

PLATE XII.

*Palinostus lalandii* (p. 86).

(*Palinurus lalandii* on Plate.)

Fig. 1. Lateral view, with part of the carapace removed to show the arrangement of the branchiæ.

„ 1*h*. First gnathopod.

„ 1*i*. Second gnathopod.

*Panulirus penicillatus* (p. 82).

„ 2. Lateral view of pereion, with part of the carapace removed to show the arrangement of the branchiæ; slightly enlarged.

„ 2*h*. First gnathopod.

„ 2*i*. Second gnathopod.





PLATE XII.

PLATE XIIA.

PALINURIDÆ—Development.

- Fig. 1. Brephalos of *Palinurus vulgaris*, extracted from the ovum; magnified twenty-five times (p. 89).
- .. 2. Brephalos of a *Palinurus* (?), taken off Samboangan; magnified twenty-eight times (p. 90).
- .. 3. Phyllosoma of a *Panulirus* (?), taken off the Celebes Islands; magnified nine times (p. 91).
- .. 4. Cephalon of fig. 2; *a, a*, bases of ophthalmopoda; *b, b*, bases of first pair of antennæ; *c, c*, bases of second pair of antennæ.








PLATE XII.

PLATE XII<sub>B</sub>.

PALINURIDÆ—Development (p. 89).

Fig. 1. Phyllosoma of a *Panulirus* (?), taken off the Fiji Islands; magnified seven times (p. 91).

„ 2. Pleon of the same; *v*, posterior pleopod.

„ 3. Nervous system of a Phyllosoma of *Panulirus* (?) (p. 91).

*a,a*. Ophthalmopoda.

*b,b*. First antennæ.

*c,c*. Second antennæ.

*c'*. The phymacerite.

*g,g*. Green gland (*3gg*, shows structure enlarged).

*d*. Mandibles.

*e*. First pair of siagnopoda.

*Hp*. Hepatic ducts.

*F-H*. Ganglia of thorax.

*K-O*. Ganglia of pereion.

*o,o*. Posterior pair of pereiopoda.

*P-T*. Ganglia of pleon.

*v*. Sixth somite of pleon.

*z,r*. Rhipidura.

SCYLLARIDÆ—Development (p. 95).

„ 4*b*. Second antenna of a Phyllosoma of one of the Scyllaridæ (p. 96).


„ 4*e*. First siagnopod.

„ 4*f*. Second siagnopod.

„ 4*g*. First gnathopod (?).

„ 4*h*. Second gnathopod (?).

„ 4*q*. Podobranchia and mastigobranchia of pereiopod: not yet escaped from the embryonic sac.



1. 2. 3. 4. 5. 6. 7. 8. 9. 10. 11. 12. 13. 14. 15. 16. 17. 18. 19. 20. 21. 22. 23. 24. 25. 26. 27. 28. 29. 30. 31. 32. 33. 34. 35. 36. 37. 38. 39. 40. 41. 42. 43. 44. 45. 46. 47. 48. 49. 50. 51. 52. 53. 54. 55. 56. 57. 58. 59. 60. 61. 62. 63. 64. 65. 66. 67. 68. 69. 70. 71. 72. 73. 74. 75. 76. 77. 78. 79. 80. 81. 82. 83. 84. 85. 86. 87. 88. 89. 90. 91. 92. 93. 94. 95. 96. 97. 98. 99. 100.

1. 2. 3. 4. 5. 6. 7. 8. 9. 10. 11. 12. 13. 14. 15. 16. 17. 18. 19. 20. 21. 22. 23. 24. 25. 26. 27. 28. 29. 30. 31. 32. 33. 34. 35. 36. 37. 38. 39. 40. 41. 42. 43. 44. 45. 46. 47. 48. 49. 50. 51. 52. 53. 54. 55. 56. 57. 58. 59. 60. 61. 62. 63. 64. 65. 66. 67. 68. 69. 70. 71. 72. 73. 74. 75. 76. 77. 78. 79. 80. 81. 82. 83. 84. 85. 86. 87. 88. 89. 90. 91. 92. 93. 94. 95. 96. 97. 98. 99. 100.

1. 3. PALINURIDÆ, PHYLLOSOMA. DEVELOPMENT.  
4. SCYLLARIDÆ, - - - - - D.



PLATE XIIc.

PLATE XIIc.

PALINURIDÆ and SCYLLARIDÆ—Development (p. 95).

Fig. 1. Phyllosoma of *Panulirus* (?), from the West Indies; enlarged three times (pp. 94, 97).

„ 2. Phyllosoma of one of the Scyllaridæ; enlarged eight times (p. 96).



SCUDARIDÆ PHYLLOSUMA H. J. SMITH  
I. PALINURIDÆ





PLATE XII.

PLATE XIIb.

SCYLLARIDÆ—Development

- Fig. 1. *Phyllosoma furcicaudatum*. Young of one of the Scyllaridæ; enlarged fifteen times (p. 98).
- .. 2f. Second siagnopod of *Phyllosoma verdense*; enlarged (p. 98).
- ., 2z. Telson of the same.
- ., 3p. Pleon of *Phyllosoma philippinense*; z, telson; m, posterior pair of pereopoda (p. 99).



SCYLLARIDAE

- |   |            |               |
|---|------------|---------------|
| 1 | PHYLLOSOMA | FURCICAUDATUM |
| 2 | "          | VERDENSE      |
| 3 | "          | PHILIPPINENSE |





PLATE XII<sub>E</sub>.

*Eryoncicus cæcus* (p. 122).

Fig. 1. Dorsal aspect ; enlarged five and a half times.

.. 2. Ventral aspect ; enlarged seven times.

- b.* First antenna.
- c.* Second antenna.
- k.* First pereopod.
- l.* Second pereopod.
- m.* Third pereopod.
- n.* Fourth pereopod.
- o.* Fifth pereopod.
- p.* Pleon.
- p.* First pleopod.
- z.* Telson.
- br.* Branchial plume.



ERYONEICUS CFCUS  
2 D<sup>o</sup> D<sup>o</sup> VENTRAL





PLATE XIII.

PLATE XIII.

*Polycheles crucifera* (p. 127).

Fig. 1. Dorsal aspect ; enlarged two and a half times.

- c. Anterior portion of the dorsal surface of the cephalon ; considerably enlarged, showing both pairs of antennæ (*b*, *c*), and the ophthalmopoda.
- v. Pleon ; lateral view.
- a*. Ophthalmopod seen from the front when the antennæ, *b* and *c*, are removed.
- b*. First antenna, the peduncle, viewed laterally.
- i*. Second gnathopod.
- o*. Terminal joints of the posterior pair of pereopoda.
- p*. First pleopod of male.
- q*. Second pleopod with two stylamblydes of male.





PLATE XIV.


PLATE XIV.

*Polycheles baccata* (p. 131).

- Fig. 1. Dorsal view of male ; enlarged one-fourth.
- „ 1c. Anterior portion of cephalon, showing *a*, ocular notch ; *b*, first antenna ;  
*c*, second antenna.
- „ 1p. Pleon ; lateral view.
- „ 1o, ♀. Posterior pereopod of female.
- „ 1t, oo. Posterior pair of pereopoda of male ; *pp*, first pair of pleopoda of same.
- „ 1p, ♀. First pleopod of female.
- „ 1q, ♂. Second pleopod of male, with two stylamblydes.
- „ 1q, ♀. Second pleopod of female, with one stylamblys.

*Polycheles helleri* (p. 138).

- „ 2. Dorsal view ; enlarged three times.
- „ 2c. Anterior portion of cephalon, showing *a*, ocular notch ; *b*, first antenna ;  
*c*, second antenna.
- „ 2p. Pleon ; lateral view.
- „ 2o, ♂. Posterior pereopod of male.
- „ 2p, ♂. First pleopod of male.



Ed. Bodnar, & Haworth, 1911

POLYCHELES HACCATA  
2 HELLER





PLATE XV.

PLATE XV.

*Polycheles helleri* (p. 138).

(*Pentacheles helleri* on Plate.)

- Fig. 1. Anterior portion of the dorsal surface of the cephalon, showing the ophthalmopoda and both pairs of antennæ (*b, c*).

*Pentacheles obscura* (p. 143).

- „ 2, ♀. Dorsal aspect of the female ; enlarged twice.  
„ 2P. Pleon ; lateral view.  
„ 2o, ♀. Posterior pereopod, chelate in female.


*Stereomastis suhmi* (p. 154).

(*Pentacheles suhmi* on Plate.)

- „ 3c. Dorsal view of the animal, male ; enlarged twice.  
„ 3c. Lateral half of the anterior dorsal portion of cephalon, showing ocular notch, ophthalmopod (*a*), and first antenna (*b*).  
„ 3c. First antenna.  
„ 3o, ♂. Posterior pereopod of male, showing elongation of the vas deferens.  
„ 4. Lateral view of the same animal.

*Pentacheles lævis* (p. 144).

- „ 4c. Anterior portion of cephalon ; dorsal surface, showing ocular notch with ophthalmopod.  
„ 5 ♀. Dorsal view of female ; enlarged twice.  
„ 5P. Pleon ; lateral view.  
„ 5o, ♀. Posterior pair of pleopoda in female.



1. *PENTACHELES HELLERI*  
2. " *OBSCURA*

3 & 4. *PENTACHELES SUMNERI*  
5. " *LARVA*



PLATE XVI.

PLATE XVI.

*Pentacheles gracilis* (p. 146).


Fig. 1, ♀. Dorsal view of the female ; enlarged one and three-quarter times.

- .. 1c. Part of the anterior portion of the cephalon ; enlarged, showing the ocular notch (*a*), with the position of the ophthalmopod traced beneath ; first antenna (*b*) ; second antenna (*c*).
- .. 1o. Posterior pair of pereopoda of female.
- .. 2. Lateral view of the same animal.

*Stereomastis auriculata* (p. 159).

(*Pentacheles auriculata* on Plate.)

- .. 3, ♀. Dorsal view of the female.
- .. 3c. Anterior portion of the dorsal surface of the cephalon, showing the ocular notch with the ophthalmopod (*a*).
- .. 3c''. Anterior portion of the cephalon, showing the lateral and under view, ophthalmopod (*a*), and second antenna (*c*).
- .. 3o. Chela of posterior pereopod ; female.
- .. 3q. Second pair of pleopoda.
- .. 4. Lateral view of the same animal.



1. PENTOCHELES GRACILIS ♀ (DORSAL)  
2. " " (LATERAL)

3. PENTOCHELES AURICULATA ♀ (DORSAL)  
4. " " (LATERAL)






PLATE XVII.

PLATE XVII.

*Pentacheles euthrix*, female; enlarged one and three-quarter times (p. 149).

(*Pentacheles euthrix* on Plate).

- c. Cephalon, showing ophthalmopoda (*a,a*); first pair of antennæ (*b,b*); second pair of antennæ (*c,c*).
- c''. Cephalon, one side of frontal aspect; *a*, ophthalmopoda; *b,c*, antennæ removed.
- c'''. Cephalon, under surface, showing ophthalmopod (*a*); first antenna (*b*); second antenna (*c*); phymacerite (*op*).
  - c*. Second antenna; *op*, phymacerite.
  - o'*. Fifth pereopod, under surface of terminal chela.
  - o''*. Fifth pereopod, upper surface of terminal chela.
  - P*. Pleon; lateral view.
  - p, ♀*. First pleopod.
  - q*. Second pleopod.



PENTACHTES: ENTOMO.



PLATE XVIII.

PLATE XVIII.

*Willemæsia leptodactyla* ; enlarged one and a half times (p. 163).

- v.* Pleon : lateral view.
- d.* Siagon or mandible ; from the inside.
- e.* First siagnopod or first maxilla.
- f.* Second siagnopod or second maxilla.
- g.* Third siagnopod or maxilliped.
- h.* First gnathopod.
- i.* Second gnathopod.
- k.* Coxa and basis of first pereopod, showing mastigobranchial lash and podobranchial plume.
- k''.* Chela of first pereopod.
- k'''.* Podobranchia ; enlarged.
- k''''.* Series of small plates on the inner margins of pollex and dactylos.



Willemoesia teptodactyla

Willemoesia teptodactyla

WILLEMOESIA TEPTODACTYLA





PLATE XIX.

PLATE XIX.

*Willemæsia leptodactyla* (p. 163).

- c. Cephalon, dorsal aspect. Large variety; enlarged (p. 169).
- c''. Cephalon, dorsal aspect. Type specimen.
- c'''. Inferior aspect of the anterior extremity of cephalon, showing first and second pairs of antennæ, siagon, third pair of siagnopoda, and first pair of gnathopoda, all in position.
- c'''. Metope or facial region, showing ophthalmopoda (*a,a*), position of first pair of antennæ (*b,b*), position of second pair of antennæ (*c,c*).
- b. Coxa of first antenna with ophthalmopod (*a*), in position.
- b. The same enlarged, having upper surface partially removed to show the internal structure of the auditory apparatus (*ac*).
- c. Second antenna; view from the upper surface, showing the reversed position of the extremity of the phymacerite (*ot*).
- o. Posterior pair of pereopoda of male, showing the foramen in the coxa for the passage of the vas deferens.
- p*, ♂. First pleopod of male, in relative position.
- q*, ♂. Second pleopod of male, showing two stylamblydes.
- q*, ♀. Second pleopod in female, showing one stylamblys.
- p. Pleon; lateral view.
- plc*. Internal dental apparatus at the pyloric extremity of stomach.



S. B. and nat. J. Hawlins lith.

WILLEMOESIA TEPTODACTYLA.



PLATE XX.

PLATE XX.

*Willemæsia leptodactyla* (p. 163).

Fig. 1. Pereion, with lateral wall of the carapace removed to show the position and arrangement of the branchiæ.

*c.* Metope or facial region, showing on one side the relative positions of the ophthalmopod (*a*), and first (*b*) and second antennæ (*c*), from within.

*d, d.* Siagones or mandibles (the synaphipod should have only two joints).

*d'', d''.* Metastomata.

*ma.* Cheiloglossa, as seen between the mandibles.

*m.* Third pair of pereiopoda with mastigobranchial lash and podobranchial plume.

*m'''.* One of the hairs and spines from the mastigobranchia; enlarged.

*m''.* Section of a podobranchial plume.

.. 2. Ovum enlarged, showing advanced condition of the embryo.







PLATE XXI.

PLATE XXI.

*Phoberus tenuimanus*, nat. size (p. 171).

(*Acanthacaris tenuimana* on Plate.)

- a, a.* Ophthalmopoda.
- c.* Cephalon ; *b*, first antenna, left side ; *c*, second antenna, right side.
- c.* Peduncle of second antenna, seen on the lower side.
- k.* Chela of first pereopod, left side.
- k'*. Mastigobranchial leaf of same with podobranchial plume attached.
- k''*. Section of branchial plume.
- o.* Chela of posterior pereopod.
- p.* First pleopod.
- q.* Second pleopod.
- v, z, v.* Rhipidura ; *z*, telson ; *vv*, posterior pair of pleopoda.





PLATE XXII.

PLATE XXII.

*Phoberus tenuimanus* (p. 171).

(*Acanthocaris tenuimana* on Plate.)

Pereion, with the lateral wall of the carapace removed to show the natural position and arrangement of the branchiæ; *i*, second gnathopod; *k-o*, pereiopoda.

Section of a set of branchial plumes and mastigobranchiæ; *mb*, mastigobranchiæ; *pdb*, podobranchia; *arb*, *arth*, anterior and posterior arthrobranchiæ; *plbr*, pleurobranchia. Above this figure are seen portions of the anterior and posterior arthrobranchia, showing their approximation, *in situ*.

*d*. Siagon (or mandible); from the inside.

*e*. First siagnopod.

*f*. Second siagnopod.

*g*. Third siagnopod.

*h*. First gnathopod, with basecephysis, mastigobranchial plate and podobranchial plume.

*i*. Second gnathopod, with basecephysis, mastigobranchial plate, and podobranchial plume.



ACANTHOCARIS TENUIMANA, PRANCHIA &c





PLATE XXIII.

PLATE XXIII.

*Nephropsis rosea* (p. 178).

Fig. 1. Lateral view of a specimen ; enlarged one and a half times.

„ 2. Dorsal view.

„ 1*e*. First siagnopod.

„ 1*f*. Second siagnopod.

„ 1*g*. Third siagnopod.

„ 1*h*. First gnathopod ; one of the apical spines enlarged.

„ 1*i*. Second gnathopod.

*Nephropsis submi* (p. 181).

„ 3. Lateral view of the specimen ; enlarged one and a half times.

„ 3*e*. First siagnopod.





PLATE XXIV.

PLATE XXIV.


*Nephropsis rosea* (p. 178).

Fig. 1. Ventral view ; enlarged.

- b.* First antenna.
- c.* Peduncle of the second antenna, showing the phymacerite and diagram of its extremity.
- d.* Siagon or mandible ; seen from within.
- m.* Coxa of the third pereopod, having the mastigobranchia with the podobranchial plume attached.
- p.* First pleopod, probably of male.
- q.* Second pleopod, with stylamblys.
- v.z.v.* Rhipidura ; *z*, telson ; *v,v*, posterior pair of pleopoda.

*Nephropsis suhmi* (p. 181).

,, *2v.z.v.* Rhipidura ; *z*, telson ; *v,v*, posterior pair of pleopoda.



12

1 NEPHROPSIS ROSEA.  
2 " SUHMI.





PLATE XXV.

PLATE XXV.

*Nephrops thomsoni*, nat. size (p. 185).

Fig. 1, ♂. Dorso-lateral view of the male.

.. 2, ♀. Dorso-lateral view of the female.

„ 3, *v.z.v.* Rhipidura; *vv*, sixth pair of pleopoda; *z*, telson.



18 Plate of Illustrations

- |    |          |           |            |
|----|----------|-----------|------------|
| 1. | NEPHROPS | THOMSONI. | MALE       |
| 2. | "        | "         | FEMALE     |
| 3. | "        | "         | RHIPIDURA. |



PLATE XXVI.

PLATE XXVI.

*Nephrops thomsoni* (p. 185).

Fig. 1. Carapace, with the lateral wall removed to show the branchiæ.

- d.* Siagon.
- e.* First siagnopod (basal plate probably wanting).
- f.* Second siagnopod.
- g.* Third siagnopod.
- h.* First gnathopod.
- i.* Second gnathopod.
- i''.* Mastigobranchia with podobranchial plume of same.
- i'''.* Section of the plume.
- o''.* Apex of the posterior pleurobranchial plume.
- p, ♂.* First pleopod of male, exhibiting cincinnuli.
- q.* Second pleopod with stylamblys.







PLATE XXVII.

PLATE XXVII.

*Astacopsis paramattensis* (p. 202).

Fig. 1. Dorso-lateral view, natural size.

- .. 1e. First siagnopod; 1, 2, 3, 4, homologues of the four respective joints of the typical limb—coxa, basis, ischium, and meros; spines and hairs enlarged.
- .. 1f. Second siagnopod, figures as before.
- .. 1g. Third siagnopod, figures as before; *ec*, basecephysis.
- .. 1h. First gnathopod, figures as above; 5 carpos, 6 propodos, 7 dactylos; *pb*, podobranchial plume; *arb*, arthrobranchial plume; *ec*, ecphysis.
- .. 1i. Second gnathopod, figures as before; *pb*, podobranchia; *arb*, arthrobranchia.
- .. 1m''. Podobranchia of third pereopod attached to the mastigobranchial plate.
- .. 1m'''. Section of the mastigobranchial plate and podobranchia of third pereopod, with hairs from the surface of the former; magnified.

*Astacopsis sydneyensis* (p. 204).

- .. 2. Dorsal view of the specimen; enlarged twice.





PLATE XXVIII.

## PLATE XXVIII.

*Astacopsis spinifer* (p. 194).

The lateral wall of the carapace has been removed to show the branchial arrangement; podobranchiæ removed to show the plan of the branchiæ beneath; arthrobranchiæ removed from the first two pairs of pereiopoda to show their position and that of the pleurobranchiæ.

- k.* First pair of pereiopoda.
- pdb.* Podobranchiæ.
- arb.* Arthrobranchiæ.
- plb.* Pleurobranchiæ.
- mt.* Movable appendages, probably the rudiments of the foliaceous appendages that form the ventral incubatory pouch in certain genera.
- d.* Siagon or mandible; 1 coxa, 2 basis, 3 terminal joint.
- e.* First siagnopod; 1 coxa, 2 basis, 3 ischium, 4 terminal joint.
- f.* Second siagnopod; 1 coxa, which carries the mastigobranchial plate; other numerals as before.
- g.* Third siagnopod; numerals as before; *ec*, a multiarticulate ecphysis of the basis.
- h.* First gnathopod; 1, 2, 3, as before; 4 meros, 5 carpos, 6 propodos.
- i.* Second gnathopod; numerals as before; *ec*, ecphysis; *pdb*, podobranchia.
- pdb.* Podobranchia.
- arb.a.* Anterior arthrobranchia.
- arb.p.* Posterior arthrobranchia.
- plbr.* Pleurobranchia.

} Arranged in a diagrammatic section.



ASTACOPSIS SPINIFER.





PLATE XXIX.

PLATE XXIX.

*Spongicola venusta* (p. 213).

- Fig. 1, ♀. Lateral view (length, 25 mm.); enlarged four times.
- „ 2*v*. Brephalos, in Zoea form (p. 216).
- „ 3*c*. Cephalon. The lateral wall of carapace removed to show branchiæ in position.
- „ *e*. First siagnopod.
- „ *f*. Second siagnopod.
- „ *g*. Third siagnopod.
- „ *h*. First gnathopod.
- „ *i*. Second gnathopod.
- „ *i''*. Coxa, basis, and part of ischium of the same enlarged; *ec*, rudimentary basecephysis; *mb*, rudimentary mastigobranchia; *arthb*, arthrobranchia.
- „ *i*. Branchiæ of same; *mb*, mastigobranchiæ; *a,p*, anterior and posterior arthrobranchiæ; *pl*, pleurobranchia; *pdb*, podobranchia.
- „ *k*. Branchiæ of first pereopod, letters as before.
- „ *l*. Branchiæ of second pereopod, letters as before.
- „ *m*. Branchiæ of third pereopod, letters as before.
- „ *n*. Branchiæ of fourth pereopod, letters as before.
- „ *o, pl*. Pleurobranchia of the fifth pereopod.
- „ *v, z, v*. Rhipidura; *v,v*, sixth pair of pleopoda; *z*, telson.





PLATE XXX.

PLATE XXX.

*Stenopus hispidus*, female (length, 45 mm.); enlarged twice (p. 211).

- e* First siagnopod.
- f*. Second siagnopod.
- g*. Third siagnopod.
- h*. First gnathopod.
- i*. Basal part of second gnathopod ; enlarged, showing basecephysis, mastigobranchia, and arthrobranchial plume.
- p*. First pleopod.
- q*. Second pleopod.
- h-o*. Diagrammatic plan of the branchial arrangement and proportions.



STENOPUS HISPIDUS





PLATE XXXI.

PLATE XXXI.

*Penæus canaliculatus*, var. *japonicus* (p. 245).

- a.* Ophthalmopod.
- b.* First antenna.
- c.* Second antenna.
- d.* Mandible or siagon.
- e.* First siagnopod or first maxilla.
- f.* Second siagnopod or second maxilla.
- g.* Third siagnopod or first maxillipede.
- h.* First gnathopod.
- i.* Second gnathopod.
- i, cc.* Section of the basecephysis of same.



PENAEUS CANALICULATUS





PLATE XXXII.

*Penæus canaliculatus* (p. 243).

Fig. 1, ♂. Male specimen; lateral view, enlarged twice.

„ 1'', ♂. Ventral surface, showing first pair of pleopoda, with petasma.

„ 2, ♀. Female specimen; lateral view.

„ 2'', ♀. Ventral surface of female, showing thelycum.

*Penæus canaliculatus*, var. *australiensis* (p. 248).

„ 3, ♀. Female specimen.

„ 3'', ♀. Ventral surface of female.

*Penæus canaliculatus*, var. *japonicus* (p. 247).

„ 4, ♀. Ventral surface of female, showing thelycum.



1. *PENEUS CANALICULATUS*  
 1. ♂ 2. ♀  
 3. ♂ 4. ♀

VENTRAL  
 VAR. JAPONICUS VENTRAL

2. *PENEUS CANALICULATUS* (VENTRAL)  
 3. ♂ VAR. *ALBIDUS*  
 4. ♀





PLATE XXXIII.

PLATE XXXIII.

*Penæus velutinus* (p. 253).

- Fig. 1.      Lateral view, enlarged twice.
- „   1''.      Ventral aspect of male; *p, p*, first pair of pleopoda, with petasma attached.
- „   1'''.      Ventral aspect of female; *p, p*, first pair of pleopoda.
- „   1z.      Telson.

*Penæus indicus* (p. 248).

- „   2.      Lateral view of male.
- „   2, *p, p*.      First pair of pleopoda, detached, with petasma connected.
- „   2*v, z, v*.      Rhipidura; *z*, telson; *v, v*, sixth pair of pleopoda.






PLATE XXXIV.

PLATE XXXIV.

*Penæus monodon* (p. 250).

Fig. 1. Lateral view.

„ 1'', ♂ . Ventral surface of male, showing petasma.

„ 1''', ♀. Ventral surface of female, showing thelycum.

*Penæus incisipes* (p. 257).

„ 2, ♂ . Lateral view of male.

„ 2'', ♂ . Ventral surface of male, showing petasma.

„ 2''', ♀. Ventral surface of female, showing thelycum.



1. *PENAEUS MONODON*

1' D<sup>o</sup> D<sup>o</sup> VENTRAL  
1'' D<sup>o</sup> D<sup>o</sup> D<sup>o</sup>

2. *PENAEUS INDICUS*

2' D<sup>o</sup> D<sup>o</sup> VENTRAL  
2'' D<sup>o</sup> D<sup>o</sup> D<sup>o</sup>





PLATE XXXV.

PLATE XXXV.

*Penæus anchoralis* (p. 258).

Fig. 1. Lateral view, enlarged twice.

„ 1". Ventral aspect of male, with the first pair of pleopoda and petasma attached.

„ 1"". Ventral aspect of female, showing thelycum.

„ 1z. Telson.

*Penæus philippinensis* (p. 261).

„ 2. Female specimen.

„ 2". Ventral aspect of female, with first pair of pleopoda thrown back.

„ 3. Male specimen.

„ 3". Ventral aspect of male, with first pair of pleopoda and petasma attached.



|    |                   |    |                                |
|----|-------------------|----|--------------------------------|
| 1  | PENÆUS ANCHORALIS | 2  | PENÆUS PHILIPPINENSIS          |
| 1° | D°                | 2° | D°                             |
| 1° | D°                | 2° | D°                             |
|    | VENTRAL           |    | VENTRAL                        |
|    |                   | 3  | D°                             |
|    |                   |    | D°                             |
|    |                   | 3° | PENÆUS PHILIPPINENSIS, VENTRAL |



PLATE XXXVI.


PLATE XXXVI.

*Penæus fissurus* (p. 263).

- Fig. 1. Female specimen.  
,, 1''. Ventral aspect of female, showing thelycum.  
,, 1*p*. First pleopod, detached.  
,, 1*z*. Telson.

*Penæus rectacutus* (p. 266).

- ,, 2. Female specimen.  
,, 2''. Ventral aspect of female, showing thelycum.  
,, 2*p*. First pleopod, detached.  
,, 2*z,v*. Rhipidura, seen laterally ; *z*, telson ; *v*, sixth pleopod.



1. PENÆUS FISSURUS.  
1° D° D° (VENTRAL)

2. PENÆUS RECTACUTUS.  
2° D° D° VENTRAL





PLATE XXXVII.

PLATE XXXVII.

*Penæus serratus* (p. 268).

- Fig. 1. Lateral view.
- „ 1a. Ophthalmopod.
- „ 1b. First antenna.
- „ 1'', ♂. Ventral aspect of male, with first and second pairs of pleopoda attached and thrown back, with petasma attached to former.
- „ 1q. Part of second pair of pleopoda, detached.
- „ 1''', ♀. Ventral aspect of female.
- „ 1z. Telson.
- „ 1br. Branch of branchial plume, and secondary branch ; detached.

*Penæus canaliculatus*, var. *japonicus* (p. 245).

- „ 2br. Branch of branchial plume, and secondary branch ; detached.

*Hemipenæus speciosus* (p. 303).

- „ 3br. Branch of branchial plume, and secondary branch ; detached.



1. *PENÆUS SERRATUS*. 2. *PENÆUS CANALICULATUS*. 3. *JAPONICUS*.  
4. *HEMIPENÆUS SPECIOSUS*.



PLATE XXXVIII.

PLATE XXXVIII.

*Pelonicus pectinatus*, male (p. 279).

Fig. *o, o.* Posterior pair of pereopoda; basal portion; *p, p.*, first pair of pleopoda, with petasma expanded and united in the median line.

*z.* Telson.

*br.* Branchial plume seen in section.

*br''.* Secondary branch of same; taken from the base.

*br'''.* Secondary branch of same; taken from the apex.

*mb.* Mastigobranchi








PLATE XXXIX.

PLATE XXXIX.

*Philonicus mülleri* (p. 275).

Fig. 1. Female specimen.

„ 1". Ventral aspect of female ; *p, p*, first pair of pleopoda.

„ 2. Male specimen.

„ 2". First pair of pleopoda of male, with petasma folded and attached.

*br*. Primary branch of branchial plume.

*br''*. Secondary branch of branchial plume.



1. PHILONICUS MÜLLERI  
 1" D° D° VENTRAL  
 2. D° D° VENTRAL  
 2" D D° VENTRAL



PLATE XL.

PLATE XL.

*Artemesia longinaris* (p. 281).

Fig. 1. Lateral view.

„ 2, ♀. Ventral aspect of female.

*d.* Mandible or siagon.

*e.* First maxilla or first siagnopod.

*f.* Second maxilla or second siagnopod.

*g.* First maxillipede or third siagnopod.

*h.* First gnathopod.

*i.* Second gnathopod.

*p, p.* First pair of pleopoda of male, with petasma attached.

*br.* Branchial plume.







PLATE XLI.

(ZOOLOGICAL CHALLENGER.—PART LII.—1887.)—Fff.

PLATE XLI.

*Haliporus equalis* (p. 285).

Fig. 1. Lateral view.

„ 1'', ♂. Ventral aspect of male. First pair of pleopoda, with petasma attached and thrown back.

„ 1''', ♀. Ventral aspect of female, showing thelycum.

*br.* Branchiæ, seen in position.

*br''.* Last three somites of the pleon; the arthrobranchiæ of the third and fourth pairs of pereopoda removed, to show the peculiar ducts that connect the pleurobranchiæ with their somites; *mb*, mastigobranchia of the third and fourth somites.

*Haliporus obliquirostris* (p. 286).

„ 2. Lateral aspect.

„ 2'', ♀. Ventral aspect of female.



ESPLANADE, W. 1887

1. HALIPORUS EQUALIS

1' D<sup>o</sup> D<sup>o</sup> VENTRAL  
1'' D<sup>o</sup> D<sup>o</sup> VENTRAL

2. HALIPORUS OBLIQUIPOSTIS

2 D<sup>o</sup> D<sup>o</sup> VENTRAL



PLATE XL.

PLATE XLII.

*Haliporus curvirostris* (p. 288).

Fig. 1. Lateral view.

*Haliporus lævis* (p. 289).

„ 2. Lateral view.

*Haliporus neptunus* (p. 291).


„ 3. Lateral view.

„ 3'' ♂. Ventral aspect of male. First pair of pleopoda with petasma, thrown forwards; second pair, directed backwards.

„ 3''' ♀. Ventral surface of pereion of female, showing thelycum.

*Philonicus lucasii* (p. 277).

„ 4. Lateral aspect.



1 HALIPORUS CURVIROSTRIS. 3. HALIPORUS NEPTUNUS.  
 2 D<sup>o</sup> LAEVIS. 3<sup>o</sup> D<sup>o</sup> D<sup>o</sup> VENTRAL,  
 3<sup>o</sup> D<sup>o</sup> D<sup>o</sup> VENTRAL.  
 4 PHILONICUS LUCASH





PLATE XLIII.

PLATE XLIII.

*Sicyonia sculpta* (p. 294).

- Fig. 1. Lateral view of female ; enlarged two and a half times.  
,, 1". Rostrum.  
,, 1z. Telson.

*Sicyonia carinata* (p. 294).

- ,, 2. Lateral view of male ; enlarged twice.  
,, 3. Dorsal view of same ; enlarged twice.  
,, 3b. First antenna.  
,, 3c'. Second antenna, base of scaphocerite ; under surface.  
,, 3c''. Second antenna, base of scaphocerite ; upper surface.  
,, 3c'''. Second antenna ; margin of scaphocerite.  
,, 3br. Branchiæ ; extremity of plumes, showing structure.  
,, 3q. Second pleopod.

*Sicyonia lancifer* (p. 297).

- ,, 4. Lateral view of female ; enlarged two and a quarter times.  
,, 4'', ♀. Ventral aspect of pereion of female, showing thelycum.  
,, 4z. Telson.

*Sicyonia lævis* (p. 298).

- ,, 5. Lateral view of female ; enlarged two and a half times.



1. SICYONIA SCULPTA  
 2. D<sup>o</sup> CARINATA  
 3. D<sup>o</sup> D<sup>o</sup> (DORSAL)

4. SICYONIA LANCIFER  
 5. D<sup>o</sup> D<sup>o</sup> VENTRAL  
 6. LAEVIS



PLATE XLIV.

PLATE XLIV.

*Hemipenæus spinidorsalis* (p. 301).

Fig. 1. Lateral view ; one-half enlarged.

„ 1''. Branchiæ, seen in position.

„ 1'''. Mastigobranchia, with podobranchial plume attached.

*Hemipenæus gracilis* (p. 302).

„ 2. Lateral view ; one-half enlarged.

*Hemipenæus speciosus* (p. 303).

„ 3. Lateral view.

*Hemipenæus virilis* (p. 303).

„ 4. Lateral view.



*Stomatopoda*  
*Stomatopoda*








PLATE XLV.

PLATE XLV.

*Aristeus armatus* (p. 312).

Fig. 1. Lateral aspect of male.

„ 2. Ventral aspect of another specimen.

*p.* First pleopod.

*q.* Second pleopod.

*r.* Third pleopod.



ARISTEUS ARMATUS

D5

D5

VENTRAL V



PLATE XLVI.

PLATE XLVI.

*Aristeus armatus* (p. 312).

*d.* Mandible or siagon.

*ma.* Metastomata.

*e.* First maxilla or first siagnopod.

*f.* Second maxilla or second siagnopod.

*g.* Maxillipede or third siagnopod.

*br.* Branchial plume, seen in section.

*br''.* Secondary branch of branchial plume.

Fig. 1. *Aristeus?* (young) (p. 240).



ARISTEUS ARMATUS  
 1 D♀





PLATE XLVII.

PLATE XLVII.

PENÆIDÆ (Development).

- Fig. 1. *Aristeus* (?), in Megalopa stage (p. 241).  
,, 1*k*. First pair of pereopoda.  
,, 2. *Aristeus* (?), younger Megalopa stage (p. 239).  
,, 2*v*. Fifth somite and rhipidura.  
,, 3. *Aristeus* (?), older Megalopa stage (p. 238).  
,, 3*b*. First antenna.  
,, 3*d*. Mandible, showing attachment of muscles.  
,, 3*k*. First pereopod.






PLATE XLVIII.

PLATE XLVIII.

PENÆIDÆ (Development).

- Fig. 1.     *Penæus*; Zoea stage (p. 237).  
„ 2.       *Penæus*; Megalopa stage.   *v. v.* Sixth pair of pleopoda.  
„ 2*g.*      Third siagnopod or first maxillipede.  
„ 2*h.*      First gnathopod.  
„ 2*m.*      Third pereopod.  
„ 3*br.*      Branchiæ of a young *Penæus*.  
„ 3*v, z, v.* Rhipidura of same.  
„ 4.       A young *Penæus*.  
„ 4*m.*      Chela of third pereopod.  
      *br.*     Early condition of branchial plume.



11. 10. 1887

PENÆIDÆ DEVELOPMENT





PLATE XLIX.

PLATE XLIX.

*Hemipenæus semidentatus* (p. 305).

Fig. 1. Lateral view of female.

„ 1br. Branch of branchial plume.

*Hemipenæus tomentosus* (p. 307).

„ 2, ♀. Ventral aspect of female, showing hollow between pereopoda.

„ 3, ♀. The same hollow part, containing a gelatinous mass; from another specimen.

„ 3". Section of the inferior portion of the wall of the carapace overlying the branchial chamber.



REMIPENAEUS SEMIDENTATUS  
2 3 5  
TOMENTOSUS



PLATE L.

PLATE L.

*Hemipenæus tomentosus* (p. 307).

Lateral view of female.

*c.* Second antenna with scaphocerite (under side); *phc*, phymacerite.

*br.* Branchial plume in section.



HEMIPENÆUS TOMENTOSUS





PLATE LI.

PLATE LI.

*Aristeus rostridentatus* (p. 317).

Fig. 1. Lateral view of female.

„ 1". Ventral aspect ; *p, p*, first pair of pleopoda.

*d*. Siagon or mandible.

*br*. Branchial plume in section.

*k*<sup>5</sup>. Spine from the carpos of the first pair of pereopoda.



ARISTEUS ROSTRIDENTATUS



PLATE LII.

PLATE LII.

*Hepomadus glacialis* (p. 321).

Fig. 1. Lateral view of female.

.. 1". Ventral aspect of pereion, showing the thelycum.

*ea.* Epistoma.

*d.* Siagon or mandible.

*ma.* Metastomata.

*g.* Third siagnopod or first maxillipede.

*br.* Branchial plume in section.



HEPOMADUS GLACIALIS.





PLATE LIII.

PLATE LIII.

*Peteinura gubernata* (p. 324).

Lateral view ; enlarged eight times.

*b.* First antenna.

*i.* Second gnathopod.

*k.* First pereopod.

*l.* Second pereopod.



*P. S. Blanford, W. Parker, 1881*

PETEINURA GUBERNATA

*Blanford, 1881*



PLATE LIV.

PLATE LIV.

*Benthescicymus crenatus* (p. 329).

Lateral view.

- a.* Ophthalmopod.
- b.* First antenna ; upper surface.
- b''.* First antenna ; under surface.
- c.* Second antenna.
- c''.* Second antenna ; second joint of peduncle showing the ancecerite and base of the scaphocerite.
- d.* Siagon or mandible, with metastoma, *d''*, in position.
- e.* First siagnopod.
- f.* Second siagnopod.
- g.* Third siagnopod.
- h.* First gnathopod.
- i.* Second gnathopod.



BENTHESCYMU. CRENATUS





PLATE LV.


PLATE LV.

*Benthescymus crenatus* (branchiæ) (p. 329).

Fig. 1. Pereion, with side of the carapace removed to show the arrangement and position of the branchiæ.

„ 2. Branchial plume.

„ 3. Branch of the same ; attached to the central stalk.



BENTHOSICYMUS CRENATUS.



PLATE LVI.

PLATE LVI.

*Benthescymus iridescens* (p. 335).

Fig. 1. Outer branch of sixth pleopod ; arterial vessels, red ; venous, blue ; nerve-filaments, yellow ; muscles, brown.

„ 2. Ventral nerve-cord of same.

*Eucopia australis*.

„ 3. Ventral nerve-cord.

*Sergestes atlanticus*.

„ 4. Arrangement of muscles in the sixth somite of the pleon.



1. BENTHESICYMUS IRIDESCENS 3. EUCOPIA AUSTRALIS  
 D<sup>o</sup> J<sup>o</sup> 4. SERGESTES ATLANTICUS





PLATE LVII.

PLATE LVII.

*Benthesicymus brasiliensis* (p. 332).

Fig. 1. Lateral view.

„ 1br. Section of a branchial plume.

*Benthesicymus pleocanthus* (p. 334).

„ 2. Lateral view.

„ 2br. Section of branchial plume.

*Benthesicymus iridescens* (p. 335).

„ 3. Lateral view.

„ 3a. Ophthalmopod.

„ 3a''. Diagrammatic section of eye.

„ 3br. Branch of a branchial plume.



|    |               |             |
|----|---------------|-------------|
| 1. | BENTHESICYMUS | BRASILIANUS |
| 2. | DO            | PLEOCANTHUS |
| 3. | DO            | IRIDESCENS  |



PLATE LVIII.

PLATE LVIII.

*Benthesicymus altus* (p. 336).

Fig. 1. Lateral view.

„ 1*a*. Ophthalmopod.

„ 1*z*. Telson.

„ 1*br*. Section of branchial plume.

*Benthesicymus mollis* (p. 339).

„ 2. Lateral view.

„ 2*z*. Telson.

„ 2*br*. Section of branchial plume.

*Gennadas intermedius* (p. 343).

„ 3. Lateral aspect.

„ 3*z*. Telson.

„ 3*br*. Section of branchial plume.



1. BENTHESICYMUS ALTUS
2. D. MOLLIS.
3. GENNADAS INTERMEDIUS





PLATE LIX.

PLATE LIX.

*Gennadas parvus*, male (p. 340).

Fig. 1. Lateral view.

„ 2. Another specimen containing a parasitic (?) worm.

*i.* Second gnathopod ; with dactylos enlarged.

*k.* First pereopod.

*m.* Third pereopod ; with chela enlarged.

*p,p.* Basal joint of first pair of pleopoda showing the petasmata hooked together in the middle by small cincinnuli.

*p''.* Cincinnulus.

*q.* Appendages attached to the second pleopod.

*zv.* Telson and sixth pleopod.

*br.* Section of a branchial plume.



GENNADAS PARVUS.



PLATE LX.

PLATE LX.

*Petalidium foliaceum* (p. 349).

Fig. 1. Lateral view ; enlarged twice.

„ 2. Oral apparatus ; *d,d*, distal joints of the synaphipod of the mandibles ; *d'*, epistoma ; *d'',d''*, metastoma ; *e,e*, first pair of siagnopoda.

*d*. Siagon or mandible.

*e*. First siagnopod ; detached.

*f*. Second siagnopod ; detached.

*g*. Third siagnopod ; detached.

„ 3. Branchiæ, shown in position.

„ 4. A foliaceous branchial plate.








PLATE LXI.

PLATE LXI.

*Sergestes*—Development.

*Elaphocaris* Zöea (p. 355).

Fig. 1. Dorsal view ; enlarged thirty-two times.

„ 2z. Telson.

„ 3. Rostrum of another specimen.

*Elaphocaris crassus* (p. 362).

„ 4. Dorsal view ; enlarged fifty times.

„ 4*b*. First antenna.

„ 4*c*. Second antenna.



SERGESTES



PLATE LXII.

PLATE LXII.

*Sergestes*—Development.

*Elaphocaris dohrni* (pp. 357-360).

Dorsal view ; enlarged fifty-three times.

*b.* First antenna.

*c.* Second antenna.

*g.* Third siagnopod or first maxillipede.

*h.* First gnathopod.



SERGESTES





PLATE LXIII.

PLATE LXIII.

*Sergestes* (?)—Development.

*Platysacus crenatus* (p. 363).

Fig. 1. Dorsal view ; enlarged seventy times.

„ 2. Rostrum of carapace and of epistoma.

„ *b.* First antenna.

„ *c.* Second antenna.

„ *i.* Second gnathopod.



PLATYSAURUS  
SERGESTIDAE



PLATE LXIV.

PLATE LXIV.

*Sergestes*—Development.

*Acanthosoma brevitelsonis* (p. 367).

Fig. 1. Dorsal view ; enlarged thirty-five times.

*Acanthosoma tynitelsonis* (p. 369).

„ 2. Latero-dorsal view ; enlarged twenty times.

*Acanthosoma longitelsonis* (p. 371).

„ 3. Latero-dorsal view ; enlarged forty times.



SERGESES.

1. ACANTHOSOMA BREVITELSONIS.  
D° TYNITELSONIS.





PLATE LXV.

PLATE LXV.

*Sergestes*—Development.

*Acanthosoma dorsispinalis* (370).

Fig. 1. Lateral view; enlarged twenty-five times.

*Acanthosoma lævirostratis*<sup>1</sup> (p. 374).

„ 2. Lateral view; enlarged twenty-eight times.

*Mastigopus dorsispinalis* (p. 375).

„ 3. Lateral view; enlarged twenty times.

„ *i.k.l.* Branchiæ attached to the second gnathopod, and first and second pereopoda.

*Mastigopus tenuis* (p. 428).

„ 4. Lateral view; enlarged twenty times.

„ 4*b.* Base of peduncle of first antenna, with otolith.

„ 4*c.* Second antenna.

„ 4*l.* Third pereopod; extremity.

<sup>1</sup> *Habitat*.—North of Admiralty Islands, between Stations 221 and 222. Associated with *Acanthosoma dorsispinalis* and *Mastigopus dorsispinalis*. Length, 3 mm. (0.12 in.).





PLATE LXVI.

PLATE LXVI.

*Sergestes*—Development.

*Acanthosoma macrotelsonis* (p. 373).

Fig. 1. Dorsal view ; enlarged forty times.

*Mastigopus suhmi* (p. 378).

„ 2. Dorsal view ; enlarged twenty-five times.



SERGESTES





PLATE LXVII.

PLATE LXVII.

*Sergestes semiarmis* (p. 423).

- Fig. 1. Lateral view ; enlarged eight times.  
,, 1i. Second gnathopod ; distal extremity.  
,, 1l. Second pereopod ; chela.  
,, 1m. Third pereopod ; chela.

*Sergestes semiarmis*, var. (p. 424).

- ,, 2c. Second antenna ; scaphocerite.  
,, 2k. First pereopod ; distal extremity.  
,, 2l. Second pereopod ; chela.  
,, 2n, o. Fourth and fifth pereopoda.  
,, 2z. Telson.

*Sergestes laeviventralis* (p. 425).

- ,, 3. Lateral view ; enlarged thirteen times.  
,, 3l. Second pereopod ; chela.  
,, 3n, o. Fourth and fifth pereopoda.

*Mastigopus spiniventralis* (p. 379).

- ,, 4. Lateral view ; enlarged twelve times.  
,, 4z. Telson.

*Sergestes spiniventralis* (p. 426).

- ,, 5a. Ophthalmopod and rostrum.  
,, 5l. Second pereopod ; chela.  
,, 5v. Sixth pleopod ; outer branch.

*Sergestes spiniventralis*,<sup>1</sup> var.

- ,, 6a. Ophthalmopod ; b, first antenna ; c, second antenna.  
,, 6n, o. Fourth and fifth pereopoda.

<sup>1</sup> *Habitat*.—West Pacific.



|   |                     |   |                              |
|---|---------------------|---|------------------------------|
| 1 | SERGESEUS SEMIARMUS | 2 | MALACODIPLOUS SEMIIVENTRATUS |
| 2 | 2"                  | 3 | SERGESEUS                    |
| 3 | 0"                  | 4 | 0"                           |
|   | LEVIVENTRATUS       |   |                              |



PLATE LXVIII.

PLATE LXVIII.

*Sergestes atlanticus*, male (p. 389).

Lateral view ; enlarged twice.

*b* ♂ . First antenna of male.

*b* ♀ . First antenna of female ; flagella.

*c* . Second antenna ; scaphocerite and terminal joints of peduncle.

*d* . Mandible.

*e* . First siagnopod.

*f* . Second siagnopod

*g* . Third siagnopod.



SERGESTES ATLANTICUS.





PLATE LXIX.

PLATE LXIX.

*Sergestes atlanticus* (p. 393).

*p.p.* First pair of pleopoda of male, with petasmata united in the median line.

*q ♂*. Second pleopod ; appendages attached to the base of the flagella.

*q ♀*. Second pleopod ; appendage attached to the base of the flagella.

*v*. Sixth pleopod.

*z*. Telson.





PLATE LXX.

PLATE LXX.

*Sergestes japonicus* (p. 387).

Fig. 1. Lateral view ; enlarged twice.

„ 1*a*, *a*. Ophthalmopoda, and *b*, *b*, first pair of antennæ ; *c*, *c*, scaphocerites in position, dorsal view.

„ 1*b*. First antenna.

„ 1*h*. Second gnathopod.

„ 2. Branchiæ ; seen in natural position.

*Sergestes kröyeri* (p. 388).

„ 3. Lateral view ; enlarged twice.

„ 3*a*. Ophthalmopod in position, with first antenna.

„ 4. Branchiæ ; in natural position.



SERGESTES JAPONICUS  
D<sup>o</sup> KROYER





PLATE LXXI.

PLATE LXXI.

*Sergestes prehensilis* (p. 385).

Fig. 1. Lateral view ; enlarged twice.

*a.* Ophthalmopod

*b.* First antenna.

*d.* Mandible or siagon.

*e.* First siagnopod.

*f.* Second siagnopod.

*g.* Third siagnopod.

*h.* First pereopod ; showing the prehensile character of the ultimate articulation.

P. Pereion detached, with gnathopoda *h, i*, and pereopoda *k, l, m, n, o*, connected, the larger branchial plumes removed to show the position of their attachment to the somite and the rudimentary form of the pleurobranchia.





PLATE LXXII.

PLATE LXXII.

*Sergestes dorsispinalis* (p. 394).

Fig. 1. Lateral view ; enlarged fifteen times.

„ 1c. Second antenna ; otolith.

*Sergestes nasidentatus* (p. 398).

„ 2. Lateral aspect ; enlarged twelve times.

„ 2k. First pereopod ; ultimate articulation.

„ 2l. Second pereopod ; chela.

*Sergestes diapontius* (p. 399).

„ 3. Lateral view ; enlarged six times.

„ 3i. Second gnathopod ; ultimate joints.

„ 3k. First pereopod ; ultimate articulation.

„ 3l. Second pereopod ; chela.



1 SERPENTES DORVILLEA  
2 - OF - NABIDENTAT  
3 - OF - DIABENTIS





PLATE LXXIII.

PLATE LXXIII.

*Sergestes armatus* (p. 401).

- Fig. 1. Lateral aspect ; enlarged nine times.  
,, 1*b*. First antenna.  
,, 1*z*. Telson.

*Sergestes edwardsii* (p. 403).

- ,, 2. Lateral view ; enlarged twelve times  
,, 2*c*. Second antenna ; scaphocerite.  
,, 2*k*. First pereopod.  
,, 2*br*. Branchial plume.  
,, 2*br''*. Branch of branchial plume.

*Sergestes rinkii* (p. 404).

- ,, 3. Lateral view ; enlarged twelve times.  
,, 3*z*. Telson.





PLATE LXXIV.

PLATE LXXIV.

*Sergestes oculatus* (p. 406).

Fig. 1. Lateral view ; enlarged twenty times.

.. 1*b*. First antenna.

.. 1*c*. Second antenna ; scaphocerite.

.. 1*c*". Extremity of scaphocerite.

*Sergestes ovatoculus* (p. 408).

.. 2. Lateral view ; enlarged ten times.

.. 2*z*. Telson ; extremity.

*Sergestes parvidens*(p. 409).

.. 3. Lateral aspect ; enlarged fifteen times.

.. 3*b*. First antenna ; flagella.

.. 3*c*. Second antenna ; extremity of scaphocerite.

.. 3*h*. First pereopod ; ultimate articulation.



SHRIMP  
LATERAL VIEW  
DORSAL VIEW  
VENTRAL VIEW





PLATE LXXV.

PLATE LXXV.

*Sergestes corniculum* (p. 410).

Fig. 1. Lateral view ; enlarged nine times.

„ 1z. Telson.

*Sergestes ancylops* (p. 413).

„ 2. Lateral view ; enlarged thirty times.

*Sergestes longirostris* (p. 415).

„ 3. Lateral view ; enlarged fifteen times.

„ 3z. Telson.



FIGURE 100. *Stomatopoda*.  
1. *Stomatopoda* sp.  
2. *Stomatopoda* sp.  
3. *Stomatopoda* sp.



PLATE LXXVI.

PLATE LXXVI.

*Sergestes junceus* (p. 416).

Fig. 1. Lateral view ; enlarged twenty times.

*Sergestes longispinus* (p. 417).

„ 2. Lateral view ; enlarged sixteen times.

„ 2*b*. First antenna ; outer flagellum.

„ 2*c*. Second antenna ; extremity of scaphocerite.

„ 2*k*. First pereopod ; distal joints.

*Sergestes penerinkii* (p. 418).

„ 3. Enlarged sixteen times.

*Sergestes fermerinkii* (p. 419).

„ 4. Enlarged twenty-three times.



1. *LENGSTRA JONGERUS*  
 2. DO *JONGERUS*  
 3. DO *JONGERUS*  
 4. DO *JONGERUS*





PLATE LXXVII.

PLATE LXXVII.

*Sergestes longicollus* (p. 421).

Fig. 1. Lateral view ; enlarged ten times.

*Sergestes præcollus* (p. 423).

„ 2. Lateral aspect ; enlarged ten times.

„ 2*m*. Third pereopod ; chela.



1. STREPTOS LUNULATUS  
2. " " PRORHINUS



PLATE LXXVIII.

PLATE LXXVIII.

*Sciacaris telsonis* (p. 438).

Fig. 1. Ventro-lateral aspect ; enlarged twenty times.

„ 1*m*. Third pereopod ; chela.

„ 1*p*. First pleopod ; appendage at base of flagellum.

„ 1*z*. Telson ; dorsal view.

„ 2. Mastigopus stage of same ; enlarged eighteen times (p. 439).

„ 2*c*. Second antenna ; extremity of scaphocerite.

„ 2*z*. Telson ; dorsal view.

„ 3. Acanthosoma stage of same ; enlarged twenty-one times (p. 441).

„ 3*z*. Telson ; side view.



1. *Belostomatidae*

2. *Belostomatidae*

3. *Belostomatidae*





PLATE LXXIX.

PLATE LXXIX.

Development and Structure of *Lucifer* (p. 452).

Fig. 1. Zoea of *Lucifer*; enlarged sixty times.

- b.* First antenna.
- c.* Second antenna; *c''*, scaphocerite and one hair magnified.
- e.* First siagnopod or first maxilla.
- f.* Second siagnopod or second maxilla.
- g.* Third siagnopod or maxillipede.
- h.* First gnathopod.
- i.* Second gnathopod.
- k.* First pereopod.
- l.* Second pereopod.
- m.* Third pereopod; *m''*, chela of same.





PLATE LXXX.

PLATE LXXX.

*Lucifer*—Structure of male.

Fig. 1. Lateral view, showing vas deferens on left side, containing ripe spermatophore.

„ 2. Lateral view, showing vas deferens on right side.

*ptm.* Petasma.

*c.* Cephalon, anterior portion ; showing the convolution of the green gland.

*d.* Mandible.

*m.* Third pereopod ; chela.

*q.* Second pleopod.







PLATE LXXXI.

PLATE LXXXI.

Organs of generation of *Lucifer* (p. 444).

- Fig. 1 ♀ . Female, showing ovaries with spermatophore inserted into the oviduct ; also neural cord and muscular arrangement.
- .. 2 ♀ . Female with ovum approaching extrusion, with spermatophore inserted.
- .. 3 ♂ . Male, showing spermatophore ready for expulsion. Testes in outline.
- .. 4 ♂ . Male, showing testes in position, with spermatophore in outline. Another in the process of formation.





PLATE LXXXII.

PLATE LXXXII.

*Lucifer*—Development of male (p. 463).

Figs. 1-4. Males of different ages.







PLATE LXXVIII.

PLATE LXXXIII.

*Lucifer typus.*

- Fig. 1. Lateral view of male (p. 464). Enlarged twenty-two times.
- „ 2. Lateral view of female (p. 466). Enlarged twenty-two times.
- „ 3. Cephalon, showing ophthalmopod and antennæ; the second pair with the long scaphocerite and styliform phymacerite.
- m.* Terminal extremity or minute chela of the third pair of pereopoda.



LUCIFER TYPE



PLATE LXXXIV.

PLATE LXXXIV.

*Lucifer reynaudii.*

- Fig. 1. Lateral view of male (p. 466). Enlarged twenty-two times.
- „ 2. Lateral view of female (p. 467). Enlarged twenty-two times.
- „ 3. Cephalon, showing the cerebral ganglion and the neural branch leading to the second pair of antennæ.
- „ 4. Cephalon, showing the cerebral ganglion and neural cord, also the green gland and its long winding duct connecting it with the second pair of antennæ.
- „ 5. Male, posterior portion of the sixth somite of the pleon, sixth pair of pleopoda and telson, showing the arrangement of the muscles and position of the sixth pleonic gland.
- m.* Chela of the third pair of pereopoda.







PLATE LXXXV.

PLATE LXXXV.

*Acetes indicus* (p. 442).

Fig. 1. Lateral view ; enlarged twice.

*Lucifer* (young) (p. 457).

„ 2. Lateral view.

*Zoontocaris galathea* (p. 474).

„ 3. Dorsal view ; enlarged thirty times.

*Zoontocaris approximus* (p. 475).

„ 4. Dorsal view ; enlarged twenty-four times.

*Sestertius duplicidentis* (p. 477).

„ 5. Lateral view ; enlarged eighteen times.

„ 5*c*. Second antenna.

„ 5*g*. Third siagnopod.

„ 5*h*. First gnathopod.

„ 5*i*. Second gnathopod.

„ 5*k*–5*o*. Pereiopoda.

„ 5*z*. Telson.



1. ACETES INDICUS. 3. ZOONTOCARIS GALATHEÆ  
 2. LUCIFER. 4. D9 APPROXIMUS.  
 5. SESTERTIUS DUPLICIDENTES.



PLATE LXXXVI.

PLATE LXXXVI.

*Crangon affinis* (p. 484).

Fig. 1. Carapace, dorsal view ; enlarged twice.

„ 1*b*. First antenna.

„ 1*d*. Mandible ; *e*, first siagnopod ; *f*, second siagnopod ; shown in their natural position.

„ 2. Branchiæ, in natural position.

„ 3. Rhipidura.

*Crangon vulgaris* (p. 483).

„ 4. Brephalos, lateral view.

„ 4*z*. Telson of same.

*Pontocaris propensalata* (p. 496).

„ 5. Branchiæ, in natural position.



1. CRABON AFFINIS

2. D<sup>o</sup> D<sup>o</sup> BRANCHIAE

3. CRABON AFFINIS (LATERAL VIEW)

4. D<sup>o</sup> VENTRAL VIEW

5. PONTICARIS PROPENSALAT: BRANCHIAE





PLATE LXXXVII.

PLATE LXXXVII.

*Pontophilus gracilis* (p. 487).

- Fig. 1. Ventral aspect ; enlarged four times.  
„ 2. Dorsal view ; enlarged four times.  
„ 3. Branchiæ, in natural position ; *br*, section of plume.  
*p* ♂ . First pleopod of male.  
*p* ♀ . First pleopod of female.  
*q* ♂ . Second pleopod of male.  
*q* ♀ . Second pleopod of female.





PLATE LXXXVIII.

PLATE LXXXVIII.

*Pontophilus profundus* (p. 490).

- Fig. 1.    Lateral view ; enlarged four times.
- „ 1*a*.    Section of the ophthalmus, showing the form of the ocular facets.
- „ 1*i*.    Second gnathopod.
- „ 1*k*.    First pereopod ; chela.
- „ 1*l*.    Second pereopod ; chela.
- „ 1*p* <sup>†</sup>. First pleopod of male.
- „ 1*z*.    Telson.

*Pontophilus junceus* (p. 491).

- „ 2.    Lateral view; enlarged twice.
- „ 3.    Dorsal view of carapace.
- „ 4.    Rhipidura.







PLATE LXXXIX.

PLATE LXXXIX.

*Crangon vulgaris* (p. 483).

Fig. 1. Structure of the pereionic viscera of the brephalos; *a*, ophthalmus; *b*, first joint of first antenna; *c*, first joint of second antenna; *d*, mandible; *e*, first siagnopod; *f*, second siagnopod; *g*, third siagnopod; *h*, first gnathopod; *i*, second gnathopod; *k*, *l*, *m*, *n*, buds of four successive pereiopoda; *gc*, stomach, the outer double line showing its area when dilated, the inner or convoluted double line showing its contour in rhythmical contraction; *cd*, heart.

„ 1*a*“. Lenses of the ophthalmus, showing the numerical increase during development.

*Sabinea septemcarinata* (p. 493).

„ 2. Dorsal view; enlarged twice.

*Parathanas decorticus* (p. 530).

„ 3. Lateral view; enlarged twelve times.

„ 3*c*. Second antenna; scaphocerite as seen within its exuvium.

„ 3*k*–3*n*. First four pairs of pereiopoda within their exuvium.

„ 3*v*. Fifth pleopod; portion of outer branch within its exuvium.

„ 3*v*“. Fifth pleopod; one of the hairs enlarged.

*Alpheus* (Brephalos) (p. 538).

„ 4. Lateral view. From a specimen of *Alpheus* hatched by Dr. Power.

„ 4*b*. First antenna.

„ 4*c*. Second antenna.

„ 4*z*. Telson.

*Latreutes planus* (p. 584).

„ 5. Lateral view; enlarged twelve times.

*Latreutes unidentatus* (p. 586).

„ 6. Lateral view; enlarged twelve times.



1. CRANGON. . . . . 4 ALPHEUS .  
 2. SABINEA SEPTEMCARINATA. 5 LATREUTES PLANUS  
 3. PARATHANAS DECORTICUS. 6. D° UNIDENTATUS.



PLATE XC.

PLATE XC.

*Sabinea septemcarinata* (p. 493).

- Fig. 1.    Ventral surface of pereion.  
,, 1*a*.    Ophthalmopoda ; *b*, first pair of antennæ.  
,, 1*l*.    Second pereiopod.  
,, 1*p* ♂ . First pleopod of male.  
,, 1*p* ♀ . First pleopod of female.  
,, 1*z*.    Telson ; *v*, sixth pleopod.

*Pontocaris propensalata* (p. 496).

- ,, 2.    Dorsal view ; enlarged twice.  
,, 2*b*.    First antenna.  
,, 2*c*.    Second antenna ; part of flagellum enlarged.  
,, 2*i*.    Second gnathopod.  
,, 2*k*.    First pereiopod.  
,, 2*k''*.    First pereiopod ; chela, enlarged.  
,, 2*p*.    First pleopod.  
,, 2*q*.    Second pleopod.  
,, 3.    Ventral surface of pereion ; *k*, *l*, *m*, *n*, *o*, first to fifth pairs of pereiopoda.







PLATE XCI.

PLATE XCI.

*Pontocaris pennata* (p. 499).

Fig. 1. Dorsal view ; enlarged twice.

„ 2. Ventral aspect ; enlarged four times.

*d.* Mandible ; *d'*, upper surface of molar process ; *d''*, lower surface of molar process.

*g.* Third siagnopod.

*h.* First gnathopod.

*i.* Second gnathopod.

*k.* First pereopod ; *k''*, anterior margin of the palm.

*l.* Second pereopod.

*m.* Third pereopod.

*p.* First pleopod.

*q.* Second pleopod.






PLATE XCH.

PLATE XCII.

*Glyphocrangon granulosis* (p. 507).

Fig. 1. Dorsal aspect.

„ 2. Lateral view.

„ 3. Oral apparatus.

*b* ♂ . First antenna of male.

*b* ♀ . First antenna of female ; *b''* ♀ , first joint of same showing the acoustic aperture.

*c* ♀ . Second antenna of female.

*d* . Mandible.

*e* . First siagnopod.

*f* . Second siagnopod.

*g* . Third siagnopod.

*h* . First gnathopod.

*i* . Second gnathopod.

*k* . First pereopod.

*l* . Second pereopod.


*p* ♂ . First pleopod of male ; *p''* , margin of inner ramus.

*p* ♀ . First pleopod of female.

*q* ♂ . Second pleopod of male.

*q* ♀ . Second pleopod of female.

„ 4. Brephalos (p. 506).



*EUPHYAS RANSONII*

FIGURE 1  
FIGURE 2  
FIGURE 3  
FIGURE 4





PLATE XCIII.

PLATE XCIII.

*Glyphocrangon granulosus* (p. 507).

Fig. 1. Branchiæ, in natural position (p. 506).

„ 1br. Section of branchial plume.

*Glyphocrangon podager* (p. 516).

„ 2. Dorso-lateral view.

„ 2m. Third pereopod ; dactylos.

„ 2o. Fifth pereopod ; dactylos.

*Glyphocrangon regalis* (p. 517).

„ 3. Dorso-lateral view.

„ 3m. Third pereopod ; dactylos.

„ 3o. Fifth pereopod ; dactylos.

„ 4. Dorso-lateral view of a smooth variety.

„ 4m. Third pereopod ; dactylos.

„ 4o. Fifth pereopod ; dactylos.

*Glyphocrangon hastacauda* (p. 519).

„ 5. Dorso-lateral view.

„ 5m. Third pereopod ; dactylos.

„ 5o. Fifth pereopod ; dactylos.



1. GLYPHOCRANGON GRANULOSUS BRANCHI

DE PJDAGER

3. GLYPHOCRANGON REGAL

4. DE

DE SMOUT & H

5. GLYPHOCRANGON PASTACANOA



PLATE XCIV.

PLATE XCIV.

*Glyphocrangon aculeata* (p. 521).

Fig. 1. Dorsal view.


- „ 1*m*. Third pereopod ; dactylos.
- „ 1*o*. Fifth pereopod ; dactylos.
- „ 1*gc*. Gastric spines ; *gc'*, one of the spines isolated.

*Glyphocrangon acuminata* (p. 522).

- „ 2 ♂ . Lateral view of male.
- „ 2*m*. Third pereopod ; dactylos.
- „ 2*o*. Fifth pereopod ; dactylos.
- „ 3 ♀ . Lateral view of female.
- „ 3*m*. Third pereopod ; dactylos.
- „ 3*o*. Fifth pereopod ; dactylos.
- „ 3*p* ♂ . First pleopod of male.
- „ 3*p* ♀ . First pleopod of female ; ova attached.

*Glyphocrangon rimapes* (p. 523).

- „ 4. Dorso-lateral view.
- „ 4*m*. Third pereopod ; dactylos.
- „ 4*o*. Fifth pereopod ; dactylos.



|   |               |                          |
|---|---------------|--------------------------|
| 1 | CLYPHOCRANGON | ACULEATA                 |
| 2 | 3             | D <sup>o</sup> ACUMINATA |
| 4 |               | D <sup>o</sup> RIMAPES   |





PLATE XCV.

PLATE XCV.

*Nika processa* (p. 527).

Fig. 1. Lateral view ; enlarged three and a half times.

*b.* First antenna.

*c.* Second antenna, scaphocerite.

*d.* Mandible.

*e.* First siagnopod.

*f.* Second siagnopod.

*g.* Third siagnopod.

*h.* First gnathopod.

*i.* Second gnathopod.

*l.* Second pereopod of right side.

*ov.* Ovum.





PLATE XCVI.

PLATE XCVI.

*Athanas veloculus* (p. 529).

Fig. 1. Lateral view ; enlarged fifteen times.

*Cheirothrix parvimanus* (p. 533).

„ 2. Lateral view ; enlarged six times.

„ 2*a*, *a*. Ophthalmopoda.

„ 2*b*. First antenna.

„ 2*c*. Second antenna.

„ 2*h*. First gnathopod.

„ 2*i*. Second gnathopod.

„ 2*l*. Second pereopod.

„ 2*l'*. Second pereopod ; enlarged.

„ 2*l''*. Second pereopod ; chela, more enlarged.

„ 2*l'''*. Second pereopod ; one hair, magnified.



1 ATHANAS VELOCULUS  
2 CHEIROTHRIX PARVIMANUS





PLATE XC VII.

PLATE XCVII.

*Alpheus edwardsii* (p. 542).

Fig. 1. Lateral view ; enlarged three times.

„ 1*d*. Mandible.

„ 1*h*. First gnathopod.

„ 1*i*. Second gnathopod.

„ 1*k*. First pereopod ; larger chela.

„ 1*m*. Third pereopod ; terminal joints.

„ 1*mb*. Mastigobranchia.

*Alpheus acuto-femoratus* (p. 545).

„ 2. Lateral view ; enlarged three times.

„ 2*k*. First pereopod ; larger chela.

*Alpheus cristidigitus* (p. 546).

„ 3. Lateral view ; enlarged four times.

„ 3*k*. First pereopod ; smaller chela.

*Alpheus megacheles* (p. 547).

„ 4. First pereopod ; larger chela.



1 ALPHEUS EDWARDSII 2 ALPHEUS ADLITI FEMORATUS  
3 ALPHEUS CRISTIDIGITUS  
4 MEGACHELES



PLATE XCVIII.

PLATE XCVIII.

*Alpheus leviusculus*, var. (p. 549).

Fig. 1. Lateral view ; enlarged three times.

„ 1c. Cephalon ; *b*, first antenna ; *c*, second antenna.

„ 1*k*. First pereiopod ; left side.

„ 1*k''*. First pereiopod ; left side, pollex.

„ 1*m*. Third pereiopod ; terminal joints.

„ 1*z*. Telson.

*Alpheus crinitus* (p. 548).

„ 2. Lateral view ; enlarged four times.

„ 2c. Cephalon with first and second antennæ ; left side.

*Alpheus bermudensis* (p. 547).

„ 3. Lateral view ; enlarged three times.

„ 3*k*. First pereiopod ; larger chela.

*Alpheus longimanus* (p. 551).


„ 4. Lateral view ; enlarged three times.

„ 4c. Cephalon ; frontal margin.

„ 4*c*. Second antenna ; scaphocerite.

„ 4*k*. First pereiopod ; larger chela.

„ 4*k'*. First pereiopod ; smaller chela of left side.



1 ALPHEUS LEVIUSCULUS.  
2. " CRINITUS

3 ALPHEUS BERMUDENSIS  
4. " LONGIMANUS





PLATE XCIX.

PLATE XCIX.

*Alpheus rapax* (p. 552).

- Fig. 1. Lateral view ; enlarged twice.
- „ 1c. Cephalon ; *b*, first antenna, left side ; *c*, second antenna, right side.
- „ 1*k*. First pereopod ; larger chela.
- „ 1*k*'. First pereopod ; showing dactylos and pollex, enlarged.
- „ 1*z*. Telson.

*Alpheus crassimanus* (p. 554).

- „ 2. Lateral view ; enlarged twice.
- „ 2*k*. First pereopod ; larger chela.

*Alpheus lavis* (p. 555).

- „ 3. Lateral view ; enlarged three times.
- „ 3c. Cephalon, with first and second antennæ.
- „ 3c'. Carapace, showing vessels of circulation.
- „ 3*k*. First pereopod ; larger chela.

*Alpheus prolificus* (p. 556).

- „ 4. Lateral view ; enlarged three times.
- „ 4c. Cephalon, with first and second antennæ.
- „ 4*m*. Third pereopod.



1 ALPHEUS RAPAX

2 " CRASSIMANUS

3. ALPHEUS LAEVIS

4 PROLIFICUS



PLATE C.

PLATE C.

*Alpheus intrinsecus* (p. 557).

- Fig. 1. Lateral view ; female, enlarged three times.
- „ 1c. Cephalon, frontal region.
  - „ 1c''. Cephalon ; *b*, first antenna ; *c*, second antenna.
  - „ 1k. First pereiopod ; smaller chela.
  - „ 1q. Second pleopod, with ova attached.
  - „ 1z. Telson.

*Alpheus minus* (p. 558).

- „ 2. Lateral view ; enlarged four times.
- „ 2k. First pereiopod ; larger chela.
- „ 2k''. First pereiopod ; pollex and dactylos.

*Alpheus spiniger* (p. 560).

- „ 3. Lateral view ; enlarged three times.
- „ 3c. Cephalon, with first and second antennæ.
- „ 3k. First pereiopod ; larger chela.
- „ 3k'. First pereiopod ; smaller chela, left.
- „ 3k''. First pereiopod, showing form of dactylos.
- „ 3m. Third pereiopod ; dactylos.
- „ 3z. Telson.



1. *ASTACUS INDIANUS* 2. *ASTACUS MINOR*  
3. *ASTACUS MINOR*





PLATE CI.

PLATE CI.

*Alpheus avarus* (p. 544).

Fig. 1. Lateral view; enlarged twice.

*Alpheus neptunus* (p. 563).

- „ 2. Lateral view; enlarged twice.
- „ 2*k*. First pereopod; larger chela, left.
- „ 2*k'*. First pereopod; smaller chela, right.

*Alpheus gracilipes* (p. 561).

- „ 3. Lateral view; enlarged three times.
- „ 3*c*. Cephalon, frontal region.
- „ 3*k*. First pereopod; larger chela.
- „ 3*m*. Third pereopod; terminal joints.
- „ 3*z*. Telson.

*Alpheus biunguiculatus* (p. 562).

- „ 4. Lateral view; enlarged three times.
- „ 4*o*. Fifth pereopod; terminal joints.
- „ 4*z*. Telson.

*Betæus malleodigitus* (p. 565).

- „ 5. Lateral view; enlarged three times.
- „ 5*c*. Cephalon, frontal region.
- „ 5*c*. Second antennæ; scaphocerite.
- „ 5*k*. First pereopod; larger chela.
- „ 5*l*. Second pereopod.
- „ 5*p*. First pleopod, with parasites attached to peduncle; stylamblys detached and enlarged.
- „ 5*par*. Parasitic vesicle, detached; escaped ovum, enlarged.
- „ 5*z*. Telson.

*Betæus microstylus* (p. 566).

- „ 6. Lateral view; enlarged three times.
- „ 6*c*. Cephalon, frontal region and scaphocerites.
- „ 6*c*. Second antennæ; scaphocerite, enlarged.
- „ 6*z*. Telson.



1. ALPHEUS AVARUS  
2. " NEPTUNUS.  
3. " GRACILIPES

4. ALPHEUS BIUNGUICULATUS  
5. BETÆUS MALLEODIGITUS  
6. " MICROSTYLUS



PLATE CII.

PLATE CII.

*Paralpheus diversimanus* (p. 568).

Fig. 1. Lateral view ; enlarged three times.

*b.* First antenna.

*c.* Second antenna.

*d.* Mandible.

*e.* First siagnopod.

*f.* Second siagnopod.

*g.* Third siagnopod.

*h.* First gnathopod.

*i.* Second gnathopod.

*k.* First pereopod ; smaller chela.

*p.* First pleopod.

*q.* Second pleopod.

*r.* Third pleopod.

*vzv.* Rhipidura.



PARALPHEUS DIVERSIMANUS.





PLATE CIII.

PLATE CIII.

*Synalpheus falcatus* (p. 574).

Fig. 1. Lateral view ; enlarged three times

c. Cephalon, frontal region ; with *b*, first, and *c*, second antenna.

*d*. Mandible.

*e*. First siagnopod.

*f*. Second siagnopod.

*g*. Third siagnopod.

*h*. First gnathopod.

*i*. Second gnathopod.

*k*. First pereopod ; larger chela, left.

*k'*. Chela open to show the form of the dactylos.

*k''*. First pereopod ; smaller chela, right side.

*m*. Third pereopod ; terminal joints.

*q*. Second pleopod, bearing ova.

*zv*. Rhipidura.





PLATE CIV.

PLATE CIV.

*Latreutes ensiferus* (p. 583).

Fig 1. Lateral view; enlarged six times.

- „ 1*d*. Mandible.
- „ 1*e*. First siagnopod.
- „ 1*f*. Second siagnopod.
- „ 1*g*. Third siagnopod.
- „ 1*h*. First gnathopod.
- „ 1*i*. Second gnathopod.
- „ 1*k*. First pereopod.
- „ 1*l*. Second pereopod.
- „ 1*m*. Third pereopod; distal joints.
- „ 1*q*. Second pleopod, with ova attached.

*Platybema rugosum* (p. 579).

- „ 2. Lateral view; enlarged six times.
- „ 2*d*. Mandible.
- „ 2*h*. First gnathopod.
- „ 2*k*. First pereopod.
- „ 2*l*. Second pereopod.
- „ 2*q*. Second pleopod, with ova attached, and stylamblys detached and magnified.
- „ 2*vzv*. Rhipidura.









PLATE CV.

PLATE CV.

*Hippolyte bidentatus* (p. 591).

- Fig. 1. Lateral view of male ; enlarged three times.
- „ 1*b*. First antennæ.
  - „ 1*d*. Mandible.
  - „ 1*e*. First siagnopod.
  - „ 1*g*. Third siagnopod.
  - „ 1*h*. First gnathopod.
  - „ 1*i*. Second gnathopod.
  - „ 1*k*. First pereopod.
  - „ 1*l*. Second pereopod.
  - „ 1*m*. Third pereopod ; distal joints.
  - „ 2. Lateral view of female ; enlarged three times.
  - „ 2'. Rostrum ; magnified.

*Hippolyte projecta* (p. 594).

- „ 3. Lateral view ; enlarged three times.
- „ 3*a*. Ophthalmopod.
- „ 3*b*. First antenna.



1. Shrimp, 2. Shrimp, 3. Shrimp.  
A. B. C. D. E. F. G. H. I. J. K. L. M. N. O. P. Q. R. S. T. U. V. W. X. Y. Z.




PLATE CVI.

PLATE CVI.

*Spirontocaris spinus* (p. 596).

- Fig. 1. Rostrum, after Leach, var.  $\alpha$ .  
.. 2. Lateral view of animal, after Owen, var.  $\beta$ .  
.. 3. Lateral view of var.  $\gamma$ .  
.. 4. Lateral view of var.  $\epsilon$ .  
.. 5. Ventral aspect of same, with saccular parasite attached.  
.. 6. Ventral aspect of same, with *Phryxus* attached.  
.. 7. Lateral view of var.  $\zeta$ ; ophthalmopoda removed.  
.. 8. Rostrum of var.  $\eta$ .  
.. 9. *Phryxus*, female, ♀, with male, ♂, attached.  
.. 10. Parasite from fig. 5; enlarged.



SPIRONTOCARIS SPINUS





PLATE CVII.

PLATE CVII.

*Spirontocaris spinus*, var.  $\delta$  (p. 599).

Fig. 1. Lateral view ; enlarged three times.

- a.* Ophthalmopod.
- b.* First antenna.
- c.* Second antenna ; scaphocerite.
- d.* Mandible.
- e.* First siagnopod.
- f.* Second siagnopod.
- h.* First gnathopod.
- i.* Second gnathopod.
- k.* First pereopod.
- l.* Second pereopod.
- m.* Third pereopod ; distal joints.
- mb.* Mastigobranchia.






PLATE CVIII.

PLATE CVIII.

*Nauticaris marionis* (p. 603).

Fig. 1. Lateral view ; enlarged twice.

*a.* Ophthalmopod.

*b.* First antenna.

*c.* Second antenna, scaphocerite.

*d.* Mandible.

*e.* First siagnopod.

*f.* Second siagnopod.

*g.* Third siagnopod.

*h.* First gnathopod.

*i.* Second gnathopod.

*i''.* Second gnathopod ; mastigobranchia.

*l.* Second pereiopod.

*v.* Sixth somite ; *v''*, articulated process at the infero-posterior angle.

*vz.* Rhipidura ; lateral view.

„ 2. Branchial apparatus *in situ*.







PLATE CIX.

PLATE CIX.

*Nauticaris futilirostris* (p. 606).

Fig. 1. Lateral view ; enlarged twice.

*Hetairus gaimardii* (p. 611).

„ 2. Lateral view ; enlarged four times.

„ 2*d*. Mandible.

„ 2*g*. Third siagnopod.


„ 2*h*. First gnathopod.

*Hetairus tenuis* (p. 613).

„ 3. Lateral view ; enlarged three times.

*Hetairus debilis* (p. 615).

„ 4. Lateral view ; enlarged four times.



NAUTICARIS FUTILIROSTRIS. 3. HETAIRUS TENUIS  
 2. HETAIRUS GAIMARDII 4. OPHRYOTROCHA DEBILIS



PLATE CX.

PLATE CX.

*Nauticaris unirecedens* (p. 608).

Fig. 1. Lateral view ; enlarged five times.

*Chorismus tuberculatus* (p. 617).

„ 2. Lateral view.

„ 2*a*. Ophthalmopod.

„ 2*d*. Mandible.

*Amphiplectus depressus* (p. 623).

„ 3. Lateral view ; enlarged twice.

„ 3*d*. Mandible.


„ 3*i*. First gnathopod.

*Merhippolyte agulhasensis* (p. 619).

„ 4. Lateral view ; enlarged twice.

„ 4*a*. Ophthalmopod.

„ 4*i*. Second gnathopod.



1. NAUTICARIS UNIRECEDENS. 3. AMPHIPLECTUS DEPRESSUS.  
2. CHORISMUS TUBERCULATUS. 4. MERHIPPOLYTE AGULHASSENSIS





PLATE CXI.

PLATE CXI.

*Heterocarpus dorsalis* (p. 630).

Fig. 1 ♀. Lateral view of female.

- b.* First antenna.
- c.* Second antenna ; scaphocerite.
- d.* Mandible ; synhipod detached.
- f.* Second siagnopod.
- g.* Third siagnopod.
- h.* First gnathopod.
- i.* Second gnathopod.
- k.* First pereopod.
- l.* Second pereopod.
- m.* Third pereopod.
- p.* First pleopod.
- q.* Second pleopod.
- zv.* Rhipidura.
- ,, 2. Branchiæ in natural position.





PLATE CXII.

PLATE CXII.

*Heterocarpus alphonsi* (p. 632).

Fig. 1. Lateral view.

„ 1*l*, 1*l'*. Second pereopod ; right and left.

*Heterocarpus gibbosus* (p. 634).

„ 2. Lateral view ; enlarged twice.

„ *i*. First gnathopod.

„ *k*, *l*, *m*. First three pereopoda.

„ 2*z*. Telson.

*Heterocarpus lævigatus* (p. 636).

„ 3. Lateral view.


*Heterocarpus ensifer* (p. 638).

„ 4. Lateral view.

*Dorodotes levicarina* (p. 680).

„ 5. Lateral view ; enlarged twice.

„ 5*l*. Second pereopod.



1. HETEROCARPUS ALPHONSEI 3. HETEROCARPUS LEVIGATUS  
2. " " GIBBOSUS 4. " " ENSIFER  
5. DORODOTES LEVICARINA





PLATE CXIII.

PLATE CXIII.

*Plesionika uniproducta* (p. 641).

- Fig. 1. Lateral view.
- .. 1a, a. Ophthalmopoda.
  - .. 1b. First antenna.
  - .. 1c. Second antenna ; scaphocerite.
  - .. 1ant., 1post. Epistoma and metastomata in relative position.
  - .. 1d. Mandible.
  - .. 1e. First siagnopod.
  - .. 1f. Second siagnopod.
  - .. 1g. Third siagnopod.
  - .. 1h. First gnathopod.
  - .. 1l. Second pereopod, right.
  - .. 1l'. Second pereopod, left.

*Plesionika spinipes* (p. 646).

- .. 2. Lateral view ; enlarged twice.
- .. 2k. First pereopod, terminal joints.
- .. 2m. Third pereopod, articulation of mero-carpal joint.

*Plesionika semilævis* (p. 644).


- .. 3. Lateral view ; slightly enlarged.
- .. 3b. First antenna.

*Plesionika unidens* (p. 648).

- .. 4. Lateral view ; slightly enlarged.

*Plesionika brevirostris* (p. 650).

- .. 5. Lateral view.
- .. 5l. Second pereopod.



Sp. nat. II. 1. 1. 1. 1.

|   |                         |   |                      |
|---|-------------------------|---|----------------------|
| 1 | PLESIONIKA UNIPRODUCTA  | 3 | PLESIONIKA SEMILÆVIS |
| 2 | DO SPINIPES.            | 4 | DO UNIDENS           |
| 5 | PLESIONIKA BREVIROSTRIS |   |                      |



PLATE CXIV.

PLATE CXIV.

*Nothocaris rostricrescentis* (p. 653).

- Fig. 1. Lateral view.  
,, 1*r.c.* Rostral crest.  
,, 1*a.* Ophthalmopod.  
,, 1*b.* First antenna.  
,, 1*d.* Mandible.  
,, 1*h.* First gnathopod.  
,, 1*i.* Second gnathopod.  
,, 1*k.* First pereopod.  
,, 1*k''*. First pereopod, terminal joints.  
,, 1*z.* Telson.

*Nothocaris binoculus* (p. 656).

- ,, 2. Lateral view.  
,, 2*r.c.* Rostral crest.  
,, 2*a.* Ophthalmopod.

*Nothocaris ocellus* (p. 657).

- ,, 3. Lateral view.  
,, 3*r.c.* Rostral crest.

*Pandalus modestus* (p. 670).

- ,, 4. Lateral view; enlarged four times.  
,, 4*b.* First antenna.  
,, 4*k.* First pereopod.  
,, 4*l.* Second pereopod, right.  
,, 4*l'*. Second pereopod, left.  
,, 4*m.* Third pereopod.



1 NOTHOCARIS ROSTRICRESCENTIS 3 NOTHOCARIS OCELLUS  
2 " BINOCULUS 4 PANDALUS MODESTUS





PLATE CXV.

(ZOOLOG. CHALL. EXP.—PART LII.—1887.)—Fif.

PLATE CXV.

*Pandalus magnoculus* (p. 667).

Fig. 1. Lateral view ; enlarged twice.

„ 1*a*. Ophthalmopod.

„ 1*b*. First antenna.

„ 1*z*. Telson.

*Pandalus falcipes* (p. 668).

„ 2. Lateral view ; enlarged twice.

„ 2*a*. Ophthalmopod.

„ 2*b*. First antenna.

„ 2*z*. Telson.

*Pandalopsis amplus* (p. 671).

„ 3. Lateral view.

„ 3*a*. Ophthalmopod.

„ 3*b*. First antenna.

„ 3*i*. Second gnathopod.

„ 3*k*. First pereopod.

„ 3*z*. Telson.



1. PANDALUS MAGNOCULUS  
2. PANDALUS FALCIPES  
3. PANDALOPSIS AMPLUS



PLATE CXVI.

PLATE CXVI.

*Chlorotocus incertus* (p. 674).

Fig. 1. Lateral view ; enlarged twice.

„ 1*a*. Ophthalmopod.

„ 1*b*. First antenna.

„ 1*c*. Second antenna.

„ 1*d*. Mandible.

„ 1*e*. First siagnopod.

„ 1*f*. Second siagnopod.

„ 1*g*. Third siagnopod.

„ 1*h*. First gnathopod.

„ 1*i*. Second gnathopod.

„ 1*k*. First pereopod.

„ 2. Pereion, showing branchial arrangement.

*Dorodotes reflexus* (p. 678).

„ 3. Lateral view.

„ 3*a*. Ophthalmopod.

*Nothocaris geniculatus* (p. 661).

„ 4. Lateral view of female.

„ 4*a*. Ophthalmopod.

„ 4*q*. Second pleopod ; stylamblys detached.



2 CHLOROTOCUS INCISUS  
 JOPUODOTES REFLEXUS  
 4 NOTHOCARIS GENICULATUS





PLATE CXVII.

PLATE CXVII.

*Thalassocaris dana* (p. 683).

- Fig. 1. Lateral view ; enlarged ten times.  
,, 1*b*. First antenna.  
.. 1*c*. Second antenna ; 1*c*'', inner margin of scaphocerite, enlarged.

*Thalassocaris stimpsoni* (p. 684).

- .. 2. Dorsal view ; enlarged ten times.

*Diaphoropus versipellis* (p. 687).

- .. 3. Lateral view ; enlarged ten times.  
.. 3*k*. First pereopod.  
.. 3*l*. Second pereopod.  
.. 3*m*. Third pereopod.  
.. 3*o*. Fourth pereopod.

*Diaphoropus longidorsalis* (p. 688).

- .. 4. Lateral view ; enlarged eighteen times.



PLATE 100. CRUSTACEA.

- |                       |                            |
|-----------------------|----------------------------|
| 1. THALASSOCARIS DANE | 3. DIAPHOROPUS VERSIPELLIS |
| 2. " " STIMPSONI      | 4. " " LONGIDORSALIS       |



PLATE CXVIII.

PLATE CXVIII.

*Atya sulcatipes* (p. 694).

- Fig. 1.      Lateral view ; slightly enlarged.
- b.*      First antenna.
- c.*      Second antenna, with three articuli of the flagellum enlarged.
- d.*      Mandible.
- f.*      Second siagnopod.
- g.*      Third siagnopod.
- h.*      First gnathopod.
- i.*      Second gnathopod.
- k.*      First pereopod.
- v, z, v.* Rhipidura.



ATYA SULCATIPES





PLATE CXIX.

PLATE CXIX.

*Atya sulcatipes* (p. 694).

Fig. 1. Lateral view, showing branchiæ.

„ 1*p*. First pleopod.

„ 1*q*. Second pleopod.

*Atya serrata* (p. 699).

„ 2. Lateral view; enlarged twice.

„ 2*a*. Ophthalmopod.

*Caridina typus* (p. 704).

„ 3. Part of pereion, with appendages.

„ 3*d*. Mandible.

„ 3*i*. Second gnathopod.

„ 3*k*. First pereopod.

„ 3*l*. Second pereopod.

„ 3*m*. Third pereopod.



ATYA SULCATIPES  
D<sup>o</sup> SERRATA  
3 CARIDINA TYPUS



PLATE CXX.

PLATE CXX.

*Atya bisulcata* (p. 700).

Fig. 1. Lateral view ; enlarged twice.

*b.* First antenna.

*c.* Second antenna.

*d.* Mandible ; inner and outer surfaces.

*e.* First siagnopod.

*f.* Second siagnopod.

*g.* Third siagnopod.

*h.* First gnathopod.

*i.* Second gnathopod.

*k.* First pereopod.

*o.* Fifth pereopod.







PLATE CXXI.

PLATE CXXI.

*Kryptocaris stylofrontalis* (p. 690).

Fig. 1. Lateral view ; enlarged twelve times.

„ 1*i*. Second gnathopod.

„ 1*k*. First pereopod.

„ 1*l*. Second pereopod.

„ 1*m*. Third pereopod.

*Caricyphus cornutus* (p. 712).

„ 2. Lateral view ; enlarged twenty-four times.

„ 2*i*. Second gnathopod.

„ 2*k*. First pereopod.

*Caricyphus serramarginis* (p. 714).

„ 3. Lateral view ; enlarged twelve times.

*Caricyphus gibberosus* (p. 716).

„ 4. Lateral view ; enlarged ten times.

*Caricyphus turgidus* (p. 717).

„ 5. Lateral view ; enlarged twenty-two times.

„ 5*c*. Lateral view of rostral crest.

*Caricyphus angulatus* (p. 718).

„ 6. Lateral view ; enlarged eight times.



|   |            |                |   |            |            |
|---|------------|----------------|---|------------|------------|
| 1 | KYPTOCARIS | STYLOFRONTALIS | 4 | CARICYPHUS | GIBBEROSUS |
| 2 | CARICYPHUS | CORNATUS       | 5 | ♂          | TURGIDUS   |
| 3 | ♂          | SERRAMARGINIS  | 6 | ♂          | ANGULATUS  |



PLATE CXXII.

PLATE CXXII.

*Alpheus* (Brephalos) (p. 538).


- Fig. 1.      Megalopa stage, lateral aspect ; enlarged twenty times.
- „ 1c.      Second antenna ; 1*f*, second siagnopod ; 1*g*, third siagnopod ; 1*h*, first gnathopod ; 1*i*, second gnathopod ; 1*k*, first pereopod, right side ; 1*k'*, first pereopod, left side.

*Atya* (Brephalos) (p. 692).

- „ 2.      Ventral aspect ; enlarged twenty times.
- „ 2ov.      Ovum ; enlarged twenty times.

*Campylonotus vagans* (p. 775).

- „ 3.      Lateral view.
- „ 3b.      First antenna.
- „ 3c.      Second antenna.
- „ 3*d*, 3*d'*. Mandible ; inner and outer aspects.
- „ 3e.      First siagnopod.
- „ 3*f*.      Second siagnopod.
- „ 3*g*.      Third siagnopod.
- „ 3*h*.      First gnathopod.
- „ 3*i*.      Second gnathopod.



1. ALPHEUS BREPHALOS
2. ATYA BREPHALOS
3. CAMPTOLUNGUS LONGUS





PLATE CXXIII.

PLATE CXXIII.

*Anebocaris quadroculus* (p. 722).

Fig. 1. Lateral view ; enlarged nine times.

„ 1*k*. First pereopod.

„ 1*l*. Second pereopod.

„ 1*o*. Fifth pereopod.

*Rhomaleocaris hamulus* (p. 720).

„ 2. Lateral view ; enlarged nine times.

„ 2*c*. Rostrum.

„ 2*c*'' . Antennal tooth.

„ 2*k*. First pereopod.

*Bentheocaris exuens* (p. 724).

„ 3. Lateral view ; enlarged five times.

„ 3*c*. Rostrum ; magnified.

„ 3*a*. Ophthalmopod.

„ 3*b*. First antenna.

„ 3*c*'' . Margin of scaphocerite.

„ 3*m*. Third pereopod.

„ 3*n*. Fourth pereopod.

„ 3*o*. Fifth pereopod.

*Bentheocaris stylostratis* (p. 726).

„ 4. Lateral view ; enlarged three times.

„ 4*a*. Ophthalmopod.

„ 4*d*. Mandible.

„ 4*o*. Fifth pereopod.



CANEBOCARIS QUADRICORNIS & PENTHECARIS EXCISENS  
 CARINATOCARIS HAMILLI, & LARVAE & EGGSTAGES



PLATE CXXIV.

PLATE CXXIV.

*Pontonia meleagrinx* (p. 707).

- Fig. 1. Lateral view; enlarged three times.  
,, 2. Dorsal view.  
,, 1*b*. First antenna.  
,, 1*c*. Scaphocerite.  
,, 1*d*. Mandible.  
,, 1*e*. First siagnopod.  
,, 1*f*. Second siagnopod.  
,, 1*g*. Third siagnopod.  
,, 1*h*. First gnathopod.  
,, 1*i*. Second gnathopod.  
,, 1*p*. First pleopod.

*AcanthePHYra purpurea* (p. 733).

- ,, 3. Lateral view; enlarged twice.

*AcanthePHYra longidens* (p. 735).

- ,, 4. Lateral view; enlarged twice.  
,, 4*a*. Ophthalmopod.

*AcanthePHYra media* (p. 736).

- ,, 5. Lateral view; enlarged twice.  
,, 5*c*. Rostrum of another specimen.

*AcanthePHYra angusta* (p. 737).

- ,, 6. Lateral view.  
,, 6*a*. Ophthalmopod.

*Systellaspis lanceocaudata* (p. 758).

- ,, 7. Lateral view; enlarged a half.  
,, 7*z*. Telson.



1. 2 PONTONIA MELEAGRINÆ
- 3 ACANTHEPHYRA PURPUREA.
- 4 D° LONGIDENS.

- 5 ACANTHEPHYRA MEDIA.
- 6 D° ANGUSTA.
- 7 SYSTELLASPIS LANCEOCAUDATA.





PLATE CXXV.

PLATE CXXV.

*Acanthephyra sica* (p. 739).

Fig. 1. Lateral view of female.

- „ 1*a*. Ophthalmopod.
- „ 1*b*. First antenna.
- „ 1*c*. Second antenna; under surface of scaphocerite.
- „ 1*c''*. Upper surface of scaphocerite.
- „ 1*d*. Mandible; *ch.a*, cheiloglossa; *d*, mandible; *ma*, metastoma.
- „ 1*e*. First siagnopod.
- „ 1*f*. Second siagnopod.
- „ 1*g*. Third siagnopod.
- „ 1*i*. Second gnathopod.
- „ 1*z*. Telson.

*Acanthephyra armata* (p. 744).

- „ 2. Lateral view of male.
- „ 2*p*. First pleopod.

*Acanthephyra acanthitelsonis* (p. 745).

- „ 3. Lateral view of male.
- „ 3*b*. First antenna.
- „ 3*c*. Second antenna, showing the scaphocerite with the flagellum undergoing reproduction.
- „ 3*z*. Telson.





PLATE CXXVI.

PLATE CXXVI.

*AcanthePHYra edwardsii* (p. 747).

Fig. 1. Lateral view of male.

„ 1z. Telson.

*AcanthePHYra carinata* (p. 748).

„ 2. Lateral view of male.

„ 2a. Ophthalmopod.

„ 2z. Telson.

*AcanthePHYra acutifrons* (p. 749).

„ 3. Lateral view of male.

„ 3a. Ophthalmopod.

„ 3z. Telson.

*AcanthePHYra kingsleyi* (p. 751).

„ 4. Lateral view of male.

„ 4a. Ophthalmopod.

„ 4z. Telson.

*AcanthePHYra brevirostris* (p. 751).

„ 5. Lateral view of male.

„ 6. Lateral view of a variety (p. 752).

„ 6a. Ophthalmopod.

„ 6p. First pleopod.

„ 6z. Telson.

*AcanthePHYra brachytelsonis* (p. 733).

„ 7. Lateral view of female.

„ 7a. Ophthalmopod.

*AcanthePHYra approxima* (p. 755).

„ 8. Lateral view of male.

„ 8a. Ophthalmopod.

„ 8z. Telson.



|    |              |            |   |              |                |
|----|--------------|------------|---|--------------|----------------|
| 1. | ACANTHEPHYRA | EDWARDSII  | 5 | ACANTHEPHYRA | BREVIROSTRIS   |
| 2. | ♂            | CARINATA   | 6 | ♂            | ♂              |
| 3. | ♂            | ACUTIFRONS | 7 | ♂            | BRACHYTELSONIS |
| 4. | ♂            | REINOLDII  | 8 | ♂            | APPROXIMA      |





PLATE CXXVII.

PLATE CXXVII.

*Oplophorus typus* (p. 762).

Fig. 1. Lateral view of female ; enlarged twice.

„ 1*a*. Ophthalmopod.

„ 1*b*. First antenna.

„ 1*c*. Second antenna.

„ 1*d*. Mandible.<sup>1</sup>

„ 1*g*. Third siagnopod.

„ 1*h*. First gnathopod.

„ 1*i*. Second gnathopod.

„ 1*k*. First pereopod.

„ 1*z*. Telson.

*Oplophorus longirostris* (p. 765).

„ 2. Lateral view of female ; enlarged twice.

*Oplophorus brevirostris* (p. 766).

„ 3. Lateral view ; enlarged twice.

*Proclates biangulatus* (p. 884).

„ 4. Lateral view ; enlarged four times.

„ 4*c*. Dorsal view of carapace ; *a*, ophthalmopod ; *b*, first, and *c*, second antenna.

<sup>1</sup> The mandible is drawn from an imperfect specimen ; the synhipod is triarticulate.



|   |            |              |   |            |              |
|---|------------|--------------|---|------------|--------------|
| 1 | OPLOPHORUS | TYPUS        | 3 | OPLOPHORUS | BREVIROSTRIS |
| 2 | DO         | LONGIROSTRIS | 4 | PROCIETES  | BIANGULATUS  |



PLATE CXXVIII.

PLATE CXXVIII.

*Campylonotus semistriatus* (p. 768).

- Fig. 1. Lateral view of female ; enlarged.  
,, 2. Ventral aspect of pereion.

*Campylonotus capensis* (p. 773).

- ,, 3. Lateral view of female.

*Palæmonella orientalis* (p. 787).


- ,, 4. Lateral view ; enlarged twenty times.  
,, 4i. Second gnathopod.

*Palæmon affinis* (p. 782).

- ,, 5. Lateral view of female ; enlarged three times.  
,, 5a. Ophthalmopod.  
,, 5d. Mandible.  
,, 5i. Second gnathopod.

*Palæmon natator* (p. 784).

- ,, 6. Lateral view of female ; enlarged twice.  
,, 7. Lateral view of male ; enlarged two and a half times.



1 CAMPYLONOTUS SEMISTRIATUS  
 2 D<sup>r</sup> VENTRAL  
 3 D<sup>r</sup> CAETERIS

4 PALAMONELLA ORIENTALIS  
 5 PALAMON AFFINIS  
 6 7 D<sup>r</sup> NATATOR





PLATE CXXIX.

PLATE CXXIX.

*Bithynis lar* (p. 789).

Fig. 1. Lateral view of male.

*Bithynis grandimanus* (p. 793).

.. 2. Lateral view.

.. 2'. Second pereopod, left.

.. 3. Lateral view of variety (p. 795).

*Brachycarpus savignyi* (p. 795).

.. 4. Lateral view ; enlarged twice.

*Brachycarpus audouini* (p. 798).

.. 5. Lateral view of female ; enlarged eight times.

.. 5*k*. First pereopod, right.



10012

10013

- |                |                         |
|----------------|-------------------------|
| 1 BITHYNIS LAR | 3 BITHYNIS GRANDIMANUS  |
| 2 DO           | 4 BRACHYCARPUS FAZ GRY  |
|                | 5 BRACHYCARPUS AUDOUINI |



PLATE CXXX.

PLATE CXXX.

*Nematocarcinus undulatus* (p. 801).

Fig. 1. Lateral view of female ; enlarged half.

*d.* Mandible.

*e.* First siagnopod.

*f.* Second siagnopod.


*g.* Third siagnopod.

*h.* First gnathopod.

*i.* Second gnathopod, terminal joints.

*k.* First pereopod, terminal joints.

*q.* Second pleopod, with ova attached.



NEMATOCARCINUS UNDULATIPES





PLATE CXXXI.

PLATE CXXXI.

*Nematocarcinus lanceopes* (p. 804).

Fig. 1. Lateral view.

♀. Ventral surface of pereion of female.

♂. Ventral surface of pereion of male.

k. First pereopod ; chela.

m. Third pereopod ; coxal and basal joints, with arthrobranchial plumes and ecphysis attached.

n. Fourth pereopod ; ischio-mæral articulation, inner surface ; *n'*, outer surface.

o. Fifth pereopod ; terminal joints.

*p* ♂. First pleopod of male.

*p* ♀. First pleopod of female.

*q* ♀. Second pleopod of female, with ova attached.



NEMATOCARINUS LANCEOPES



PLATE CXXXII.

# PLATE CXXXII.

- Nothocaris spiniserratus* (p. 663).  
 Fig. 1. Lateral view; enlarged sixteen times.  
 „ 1c. Rostrum; magnified.
- Nematocarcinus longirostris* (p. 806).  
 „ 2. Carapace, lateral view.
- Nematocarcinus proximatus* (p. 808).  
 „ 3. Carapace, lateral view.
- Nematocarcinus altus* (p. 809).  
 „ 4. Carapace, lateral view.
- Nematocarcinus productus* (p. 810).  
 „ 5. Carapace, lateral view.
- Nematocarcinus tenuipes* (p. 812).  
 „ 6. Carapace, lateral view.
- Nematocarcinus parvidentatus* (p. 814).  
 „ 7. Carapace, lateral view.
- Nematocarcinus gracilis* (p. 815).  
 „ 8. Carapace, lateral view.
- Nematocarcinus paucidentatus* (p. 819).  
 „ 9. Carapace, lateral view.
- Nematocarcinus tenuirostris* (p. 817).  
 „ 10. Carapace, lateral view.
- Nematocarcinus serratus* (p. 819).  
 „ 11. Rostrum.
- Nematocarcinus hiatus* (p. 821).  
 „ 12. Carapace, lateral view.
- Nematocarcinus intermedius* (p. 821).  
 „ 13. Carapace, lateral view.
- Stochasmus exilis* (p. 823).  
 „ 14. Lateral view; enlarged three times.  
 „ 14i. Second gnathopod.



| NOTHURUS CRISTATUS |             | NOTHURUS CRISTATUS |             | NOTHURUS CRISTATUS |             |
|--------------------|-------------|--------------------|-------------|--------------------|-------------|
| NO.                | DESCRIPTION | NO.                | DESCRIPTION | NO.                | DESCRIPTION |
| 1                  | ANTENNA     | 1                  | ANTENNA     | 1                  | ANTENNA     |
| 2                  | ANTENNA     | 2                  | ANTENNA     | 2                  | ANTENNA     |
| 3                  | ANTENNA     | 3                  | ANTENNA     | 3                  | ANTENNA     |
| 4                  | ANTENNA     | 4                  | ANTENNA     | 4                  | ANTENNA     |
| 5                  | ANTENNA     | 5                  | ANTENNA     | 5                  | ANTENNA     |
| 6                  | ANTENNA     | 6                  | ANTENNA     | 6                  | ANTENNA     |
| 7                  | ANTENNA     | 7                  | ANTENNA     | 7                  | ANTENNA     |
| 8                  | ANTENNA     | 8                  | ANTENNA     | 8                  | ANTENNA     |
| 9                  | ANTENNA     | 9                  | ANTENNA     | 9                  | ANTENNA     |
| 10                 | ANTENNA     | 10                 | ANTENNA     | 10                 | ANTENNA     |
| 11                 | ANTENNA     | 11                 | ANTENNA     | 11                 | ANTENNA     |
| 12                 | ANTENNA     | 12                 | ANTENNA     | 12                 | ANTENNA     |





PLATE CXXXIII.

PLATE CXXXIII.

*Notostomus patentissimus* (p. 826).

Fig. 1. Lateral view.

*d.* Mandible.

*e.* First siagnopod.

*f.* Second siagnopod.

*g.* Third siagnopod.

*h.* First gnathopod.

*i.* Second gnathopod.

*m.* Third pereopod; with arthrobranchia, mastigobranchia, and basecephysis attached.

*o.* Fifth pereopod; terminal joints.

*p.* First pleopod.

*q.* Second pleopod.

*br.* Section of a branchial plume.





PLATE CXXXIV.

PLATE CXXXIV.

*Notostomus patentissimus* (p. 826).

Fig. 1. Ventral surface.

„ 1*a*. Ophthalmopod.

„ 1*b*. First antenna; 1*b*'', portion of flagellum; 1*b*'', section of flagellum.

„ 1*c*. Second antenna.

„ 2. Branchiæ in position.

*Notostomus murrayi* (p. 829).

„ 3. Lateral view; enlarged twice.



1. NOTOSTOMUS PATENTISSIMUS.

BRANCHIAE.





PLATE CXXXV.

(ZOOLOGICAL CHALLENGER.—PART LIII.—1888.)—FII.

PLATE CXXXV.

*Notostomus japonicus* (p. 830).

Fig. 1. Lateral view of male.

*Notostomus perlatus* (p. 831).

„ 2. Lateral view of female.

*Notostomus brevirostris* (p. 832).

„ 3. Lateral view of male.

*Notostomus longirostris* (p. 833).

„ 4. Lateral view of female.



|   |                      |   |                         |
|---|----------------------|---|-------------------------|
| 1 | NOTOSTOMUS JAPONICUS | 3 | NOTOSTOMUS BREVIROSTRIS |
| 2 | DO                   | 4 | DO                      |
|   | PERTATUS             |   | LONGIROSTRIS            |



PLATE CXXXVI.

PLATE CXXXVI.

*Tropiocaris planipes* (p. 835).

Fig. 1. Lateral view ; enlarged twice.

*Tropiocaris tenuipes* (p. 836).

„ 2. Lateral view ; enlarged twice.

*Hymenodora duplex* (p. 843).

„ 3. Lateral view ; enlarged one and a half times.

„ 3c. Second antenna ; scaphocerite.

„ 3d. Mandible.

„ 3f. Second siagnopod.

„ 3g. Third siagnopod.

„ 3h. First gnathopod.

„ 3i. Second gnathopod ; basal joints with branchiæ attached.


„ 3o. Fifth pereopod ; terminal joint.

*Hymenodora rostrata* (p. 847).

„ 4. Lateral view ; enlarged twice.

*Hymenodora mollis* (p. 841).

„ 5. Lateral view ; enlarged twice.



1. *TROPIOCARIS PLANIPES*. 3. *HYMENODORA DUPLEX*  
 2. *TROPIOCARIS TENUIPES*. 4. *HYMENODORA ROSTRATA*  
 5. *HYMENODORA MOLLIS*.





PLATE CXXXVII.

PLATE CXXXVII.

*Hymenodora glauca* (p. 847).

- Fig. 1. Lateral view ; enlarged twice.
- „ 1*d*. Mandible.
- „ 1*m.a*. Metastoma.
- „ 1*e*. First siagnopod.
- „ 1*f*. Second siagnopod.
- „ 1*g*. Third siagnopod.
- „ 1*h*. First gnathopod.
- „ 1*i*. Second gnathopod.
- „ 1*k*. First pereopod.
- „ 1*m*. Third pereopod.
- „ 1*o*. Fifth pereopod ; terminal extremity.

*Hymenodora mollicutis* (p. 848).

- „ 2. Lateral view ; enlarged a half.
- „ 2*i*. Second gnathopod.
- „ 2*k*. First pereopod ; chela.
- „ 2*br*. Section of branchia.





PLATE CXXXVIII.

PLATE CXXXVIII.

*Stylodactylus discissipes* (p. 851).


- Fig. 1. Lateral view of male ; enlarged twice.  
,, 1*c*. Second antenna ; peduncle and part of frontal margin.  
,, 1*i*. Second gnathopod ; 1*i'*, dactylos.  
,, 1*k*. First pereopod.  
,, 1*k'*. First pereopod ; articulation of chela reversed.  
,, 1*r*z*r*. Rhipidura.

*Stylodactylus orientalis* (p. 854).

- ,, 2. Branchiæ in position.  
,, 2*d*. Mandible.  
,, 2*e*. First siagnopod.  
,, 2*f*. Second siagnopod.  
,, 2*g*. Third siagnopod.

*Stylodactylus bimaxillaris* (p. 855).

- ,, 3. Lateral view of female ; enlarged twice.  
,, 3*h*. First gnathopod.



1 STYLODACTYLUS DISCISSIPES  
2 ORIENTALIS .  
3 DO BIMAXILLARIS .





PLATE CXXXIX.

PLATE CXXXIX.

*Leptochela serratorbita* (p. 859).

- Fig. 1. Lateral view ; enlarged five times.
- „ 1c. Frontal margin of carapace, showing ophthalmopod in position.
  - „ 1d. Mandible.
  - „ 1e. First siagnopod.
  - „ 1f. Second siagnopod.
  - „ 1h. First gnathopod.
  - „ 1i. Second gnathopod.
  - „ 1k. First pereopod ; chela.
  - „ 1o. Fifth pereopod.
  - „ 1z. Telson.

*Leptochela gracilis* (p. 860).

- „ 2. Lateral view of male ; enlarged.
- „ 2o. Fifth pair of pereiopoda.
- „ 2br. Branchial plume ; *br''*, section of same.

*Leptochela robusta* (p. 862).

- „ 3. Lateral view of female ; enlarged three times.
- „ 4. Lateral view of male ; enlarged three times.



- |       |            |               |
|-------|------------|---------------|
| 1.    | LEPTOCHELA | SERRATORBITA. |
| 2.    | D°         | GRACILIS.     |
| 3, 4. | D°         | ROBUSTA.      |



PLATE CXL.

PLATE CXL.

*Pasiphæa cristata* (p. 865).

Fig. 1. Lateral view of female ; enlarged a half.

*k.* First pereopod ; chela.

*k'.* First pereopod ; extremity of pollex and dactylos magnified.

*m.* Third pereopod.

*n.* Fourth pereopod.

*o.* Fifth pereopod.

*p.* First pleopod ; inner ramus in position.

*q.* Second pleopod with stylamblys.

*v.* Sixth pleopod ; portion enlarged, showing the diæresis.

*br.* Section of branchial plume.

„ 2. Branchiæ in position.

„ 3. Parasite ; 3', the same, enlarged.



PASIPHÆA CRISTATA

0°

0°

BRANCHIÆ.





PLATE CXLI.

PLATE CXLI.

*Pasiphæa cristata* (p. 865).

Fig. 1*d*. Mandible.

„ 1*e*. First siagnopod.

„ 1*f*. Second siagnopod; 1*f*'', two hairs, enlarged.

„ 1*g*. Third siagnopod.

„ 1*h*. First gnathopod.

*Pasiphæa amplidens* (p. 870).

„ 2. Lateral view of carapace.

*Pasiphæa acutifrons* (p. 871).

„ 3. Lateral view.

„ 3*z*. Telson.

*Orphania tenuimana* (p. 872).

„ 4. Lateral view; enlarged.

„ 4*k*. First pereopod.

„ 4*l*. Second pereopod.

„ 4*z*. Telson.



Dr. J. S. G. Allen

- |                       |                         |
|-----------------------|-------------------------|
| 1. PASIPHÆA CRISTATA. | 3. PASIPHÆA ACUTIFRONS. |
| 2. DO? AMPLIDENS.     | 4. ORPHANIA TENUIMANA   |



PLATE CXLII.

PLATE CXLII.

*Oodeopus geminidentatus* (p. 876).

Fig. 1. Lateral view ; enlarged twenty-five times.

„ 1z. Telson.

*Oodeopus serratus* (p. 877).

„ 2. Lateral view ; enlarged twenty-five times.

„ 2z. Telson.

*Oodeopus serratus*, var. (p. 878).

„ 3. Lateral view ; enlarged fifteen times.

*Oodeopus armatus* (p. 879).

„ 4. Lateral view ; enlarged twenty times.

*Oodeopus longispinus* (p. 881).

„ 5. Lateral view ; enlarged twenty times.

„ 5z. Telson.



1. OODEOPUS GEMINIDENTATUS    3. OODEOPUS SERRATUS, var.  
2.    D<sup>o</sup>    SERRATUS    4.    D<sup>o</sup>    ARMATUS  
5. OODEOPUS LONGISPINUS.





PLATE CXLIII.

(ZOOLOG. CHALL. EXP.—PART III.—1888.)—Fff.

PLATE CXLIH.

*Oodeopus intermedius* (p. 879).

Fig. 1. Lateral view ; enlarged twenty times.

„ 1*v.z.v.* Rhipidura.

*Oodeopus duplex* (p. 880).


„ 2. Lateral view ; enlarged twenty times.

*Oodeopus gibbosus* (p. 882).

„ 3. Lateral view ; enlarged twenty-two times.

„ 3*v.z.v.* Rhipidura.

„ 4. Dorsal view.



OODEOPUS INTERMEDIUS.  
 2. D<sup>2</sup> DUPLEX  
 3, 4. D<sup>4</sup> GIBBOSUS



PLATE CXLIV.

PLATE CXLIV.

*Isotopus arcurostris* (p. 886).

Fig. 1. Lateral view ; enlarged ten times.

„ 1*d*. Mandible.

„ 1*f*. Second siagnopod.

„ 1*g*. Third siagnopod.

„ 1*br*. Branchial plume.

„ 1*c*. Cephalon.

*Hectarthropus exilis* (p. 889).

„ 2. Lateral view ; enlarged ten times.

„ 2*t*. Fifth somite ; dorsal aspect.

*Hectarthropus compressus* (p. 891).

„ 3. Lateral view ; enlarged eleven times.


*Hectarthropus expansus* (p. 892).

„ 4. Dorsal view ; enlarged twelve times.

„ 5. Lateral view ; enlarged twelve times.

*Hectarthropus tenuis* (p. 893).

„ 6. Lateral view ; enlarged ten times.



1. ICOTOPUS ARCUOSTRIS.      3. HECTARTHROPUS COMPRESSUS.  
 2. HECTARTHROPUS EXILIS.      4, 5.      D?      EXPANSUS.  
 6 HECTARTHROPUS TENUIS.





PLATE CXLV.

(ZOOLOG. CHALL. EXP.—PART LII.—1888.)—Fff.

PLATE CXLV.

*Eretmocarís remipes* (p. 895).

Fig. 1. Lateral view ; enlarged ten times.

*Eretmocarís longicaulis* (p. 897).

„ 2. Lateral view ; enlarged eighteen times.

*Eretmocarís stylorostris* (p. 898).

„ 3. Lateral view ; enlarged fifteen times.

*Eretmocarís corniger* (p. 900).

„ 4. Lateral view ; enlarged eighteen times.



1. ERETMOCARIS REMIPES.      3. ERETMOCARIS STYLOROSTRIS.  
2.      D<sup>o</sup>      LONGICAULIS.      4.      D<sup>o</sup>      CORNIGER.




PLATE CXLVI.

PLATE CXLVI.

*Amphion*—Development (p. 902).

- Fig. 1. Zoea with two pairs of legs; ventral aspect; enlarged thirty times; *h*, first gnathopod; *i*, second gnathopod.
- „ 2z. Telson of a second specimen (pp. 903, 915).
- „ 3. Young with three pairs of legs; ventral aspect of cephalon and pereion; *ph*, *ph*, phymacerite (pp. 903, 915).
- „ 3d. Oral appendages.
- „ 4. Young with fourth pair of legs (*l*) in process of development; ventral aspect; enlarged twenty times (pp. 905, 915).
- „ 5. A second specimen; portion of pereion showing the fourth pair (*l*) biramose (p. 905).
- „ 6. Cephalon of a specimen with five pairs of legs attached to the pereion; ventral surface, magnified; *ph*, *ph*, phymacerite.



AMPHION






PLATE CXLVII.

(ZOOLOG. CHALL. EXP.—PART LII.—1888).—FII.

PLATE CXLVII.

*Amphion reynaudii* (p. 906).

- Fig. 1. Female with six pairs of legs; ventral aspect; enlarged thirteen times; *h, i*, first and second gnathopods; *k, l, m, n*, first to fourth pereopods; *n.c.* nucleated cells near the fourth pair of legs; *n.c'*, the same more magnified.
- „ 1*br.* Branchial plume.
- „ 1*cl.* Mandible.
- „ 1*e.* First siagnopod.
- „ 1*f.* Second siagnopod.
- „ 1*g.* Third siagnopod.
- „ 1*z.* Telson.
- „ 2. Cephalon and pereion of a specimen (adult male?) with seven pairs of legs; ventral aspect; enlarged sixteen times; *h, i*, first and second gnathopods; *k-o*, pereiopoda; *te*, testes (?); *te''*, structure believed by Suhm to be testes; *h.c.*, hepatic cells; *h.c''*, the same, magnified (p. 910).
- „ 2*br.* Branchial plume, enlarged.



1. AMPHION REYNAUDII.

2 D° D° ADULT



PLATE CXLVIII.

PLATE CXLVIII.

*Amphion provocatoris* (p. 913).

- Fig. 1. Ventral aspect ; enlarged seven times.  
.. 2. Cephalon of the same ; dorsal aspect.  
.. c. Tooth on dorsal surface of carapace.  
.. *p*. First pleopod of male (?).  
.. *q*. Second pleopod of same.  
.. *z*. Telson ; extremity.



Plate 200.

AMPHION PROVOCATORIS.





PLATE CXLIX.

PLATE CXLIX.

- Fig. 1. *Sylon challenger*, attached to *Spirontocaris spinus*, a part of the right side of the Shrimp having been taken away; *x*, the injured part at the surface of the sac; I., II., III., IV., and V., the first to the fifth pleopods of the left side of the Shrimp; *Va*, the fifth pleopod of the right side. Magnified 13 diameters.
- .. 2. *Sylon challenger*, seen from below, with a part of the wall of the host attached; *e*, the ovary of the parasite. Magnified 12 diameters.
- .. 3. *Sylon challenger*, front view; *x*, the injured part at the surface. Magnified 12 diameters.
- .. 4. *Sylon schneideri*, front view; *a*, openings leading into the mantle-cavity; B, transverse section of *Hippolyte pusiola*. Magnified 12 diameters.
- .. 5. *Sylon schneideri*, seen from the side, the lateral parts of the exoskeleton of the second and third abdominal segment of the *Hippolyte* being taken away; *a* and B as in fig. 4; II.-IV., second to fourth pleopod. Magnified 12 diameters.



1-3. SYLLON CHALLENGERI.

4, 5. SYLLON SCHNEIDERI



PLATE CL.

## PLATE CL.

Fig. 1. *Sylon challenger*, part of a transverse section, passing through the ovary, *e*, and the "Eikittdruse," A; *a*, connective tissue of the mantle; *b*, epithelium surrounding the visceral mass; *c*, ramified tubes of the gland; *d*, female genital pore. Magnified 136 diameters.

„ 2. Part of the mantle and visceral mass of *Sylon schneideri*; *a*, connective tissue of the mantle; *b*, epithelium surrounding the visceral mass; *e*, ovary; *f*, chitinous outer wall of the mantle; *g*, thickening of the chitinous outer wall at the place where later on the openings of the mantle-cavity occur; *h*, epithelium of the outer side of the mantle; *i*, epithelium at the inner side of the mantle; *l*, blood-plasma (?); *n*, nervous system. Magnified 136 diameters.

„ 3. Part of the visceral mass of *Sylon schneideri*; *b* and *e* as in fig. 2; *c*, connective tissue; *l*, blood-plasma (?). Magnified 136 diameters.

„ 4. Section through a part of *Hippolyte pusiola*, with *Sylon schneideri* attached to it; *e*, ovary of the parasite; *n*, nervous system of the *Hippolyte*; B, basilar membrane; *r*, roots of the *Sylon*; *m*, abdominal flexor muscles of the host. Magnified 30 diameters.

„ 5. Section in front of the attachment of *Sylon schneideri* to *Hippolyte pusiola*; *e*, *m*, *n*, and *r* as in fig. 4; *t*, testis. Magnified 30 diameters.

„ 6. Section of the same *Sylon* as in the two foregoing figures; *e*, *t*, as in fig. 5; *t'*, communication of the interior of the testis with the mantle-cavity. Magnified 41 diameters.

Fig. 2  
X 136



Fig. 3  
X 136



Fig. 4  
X 136



Fig. 5  
X 136



Fig. 6  
X 136



















SMITHSONIAN INSTITUTION LIBRARIES



3 9088 00722 8869